

ON A RAMIFICATION BOUND OF TORSION SEMI-STABLE REPRESENTATIONS OVER A LOCAL FIELD

SHIN HATTORI

ABSTRACT. Let p be a rational prime, k be a perfect field of characteristic p , $W = W(k)$ be the ring of Witt vectors, K be a finite totally ramified extension of $\text{Frac}(W)$ of degree e and r be a non-negative integer satisfying $r < p - 1$. In this paper, we prove the upper numbering ramification group $G_K^{(j)}$ for $j > u(K, r, n)$ acts trivially on the p^n -torsion semi-stable G_K -representations with Hodge-Tate weights in $\{0, \dots, r\}$, where $u(K, 0, n) = 0$, $u(K, 1, n) = 1 + e(n + 1/(p - 1))$ and $u(K, r, n) = 1 - p^{-n} + e(n + r/(p - 1))$ for $1 < r < p - 1$.

1. INTRODUCTION

Let p be a rational prime, k be a perfect field of characteristic p , $W = W(k)$ be the ring of Witt vectors and K be a finite totally ramified extension of $K_0 = \text{Frac}(W)$ of degree $e = e(K)$. We normalize the valuation v_K of K as $v_K(p) = e$ and extend this to any algebraic closure of K . Let the maximal ideal of K be denoted by m_K , an algebraic closure of K by \bar{K} and the absolute Galois group of K by $G_K = \text{Gal}(\bar{K}/K)$. Let $G_K^{(j)}$ denote the j -th upper numbering ramification group in the sense of [10]. Namely, we put $G_K^{(j)} = G_K^{j-1}$, where the latter is the upper numbering ramification group defined in [18].

Consider a proper smooth scheme X_K over K and put $X_{\bar{K}} = X_K \times_K \bar{K}$. Let $\mathcal{L} \supseteq \mathcal{L}'$ be G_K -stable \mathbb{Z}_p -lattices in the r -th étale cohomology group $H_{\text{ét}}^r(X_{\bar{K}}, \mathbb{Q}_p)$ such that the quotient \mathcal{L}/\mathcal{L}' is killed by p^n . In [10], Fontaine conjectured the upper numbering ramification group $G_K^{(j)}$ acts trivially on the G_K -modules \mathcal{L}/\mathcal{L}' and $H_{\text{ét}}^r(X_{\bar{K}}, \mathbb{Z}/p^n\mathbb{Z})$ for $j > e(n + r/(p - 1))$ if X_K has good reduction. For $e = 1$ and $r < p - 1$, this conjecture was proved independently by himself ([11], for $n = 1$) and Abrashkin ([3], for any n), using the theory of Fontaine-Laffaille ([13]) and the comparison theorem of Fontaine-Messing ([14], see also [5] and [7]) between the étale cohomology groups of X_K and the crystalline cohomology groups of the reduction of X_K . From these results, they also showed some rareness of a proper smooth

Date: June 20, 2009.

Supported by 21st Century COE program “Mathematics of Nonlinear Structure via Singularity” at Department of Mathematics, Hokkaido university, and by the JSPS International Training Program (ITP).

scheme over \mathbb{Q} with everywhere good reduction ([11, Théorème 1], [2, Section 7]). In fact, they proved this ramification bound for the torsion crystalline representations of G_K with Hodge-Tate weights in $\{0, \dots, r\}$ in the case where K is absolutely unramified.

On the other hand, for a torsion semi-stable representation with Hodge-Tate weights in the same range, a similar ramification bound for $e = 1$ and $n = 1$ is obtained by Breuil (see [7, Proposition 9.2.2.2]). He showed, assuming the Griffiths transversality which in general does not hold, that if $e = 1$ and $r < p - 1$, then the ramification group $G_K^{(j)}$ acts trivially on the mod p semi-stable representations for $j > 2 + 1/(p - 1)$.

In this paper, we prove a ramification bound for the torsion semi-stable representations of G_K with Hodge-Tate weights in $\{0, \dots, r\}$ with no assumption on e but under the assumption $r < p - 1$. Let π be a uniformizer of K , $E(u) \in W[u]$ be the Eisenstein polynomial of π over W and S be the p -adic completion of the divided power envelope of $W[u]$ with respect to the ideal $(E(u))$. Consider a category $\text{Mod}_{/S_\infty}^{r, \phi, N}$ of filtered (ϕ_r, N) -modules over the ring S and a G_K -module

$$T_{\text{st}, \underline{\pi}}^*(\mathcal{M}) = \text{Hom}_{S, \text{Fil}^r, \phi_r, N}(\mathcal{M}, \hat{A}_{\text{st}, \infty})$$

for $\mathcal{M} \in \text{Mod}_{/S_\infty}^{r, \phi, N}$, where $\hat{A}_{\text{st}, \infty}$ is a p -adic period ring ([6]). Then our main theorem is the following.

Theorem 1.1. *Let r be a non-negative integer such that $r < p - 1$. Let \mathcal{M} be an object of the category $\text{Mod}_{/S_\infty}^{r, \phi, N}$ which is killed by p^n . Then the j -th upper numbering ramification group $G_K^{(j)}$ acts trivially on the G_K -module $T_{\text{st}, \underline{\pi}}^*(\mathcal{M})$ for $j > u(K, r, n)$, where*

$$u(K, r, n) = \begin{cases} 0 & (r = 0), \\ 1 + e(n + \frac{1}{p-1}) & (r = 1), \\ 1 - \frac{1}{p^n} + e(n + \frac{r}{p-1}) & (1 < r < p - 1). \end{cases}$$

We can check that this bound is sharp for $r \leq 1$ (Remark 5.15).

From this theorem and [10, Proposition 1.3], we have the following corollary.

Corollary 1.2. *Let the notation be as in the theorem and L be the finite extension of K cut out by the G_K -module $T_{\text{st}, \underline{\pi}}^*(\mathcal{M})$. Namely, the finite extension L is defined by*

$$G_L = \text{Ker}(G_K \rightarrow \text{Aut}(T_{\text{st}, \underline{\pi}}^*(\mathcal{M}))).$$

Let $\mathfrak{D}_{L/K}$ denote the different of the extension L/K . Then we have the inequality

$$v_K(\mathfrak{D}_{L/K}) < u(K, r, n)$$

for $r > 0$ and $v_K(\mathfrak{D}_{L/K}) = 0$ for $r = 0$.

Combining these results with a theorem of Liu ([17, Theorem 2.3.5]) or a theorem of Caruso ([8, Théorème 1.1]), we will show the corollary below.

Corollary 1.3. *Let r be a non-negative integer such that $r < p - 1$. Then the same bounds as in Theorem 1.1 and Corollary 1.2 are also valid for the torsion G_K -modules of the following two cases:*

- (1) *the G_K -module \mathcal{L}/\mathcal{L}' , where $\mathcal{L} \supseteq \mathcal{L}'$ are G_K -stable \mathbb{Z}_p -lattices in a semi-stable p -adic representation V with Hodge-Tate weights in $\{0, \dots, r\}$ such that \mathcal{L}/\mathcal{L}' is killed by p^n .*
- (2) *the G_K -module $H_{\text{ét}}^r(X_{\bar{K}}, \mathbb{Z}/p^n\mathbb{Z})$, where X_K is a proper smooth algebraic variety over K which has a proper semi-stable model over \mathcal{O}_K and r satisfies $er < p - 1$ for $n = 1$ and $e(r + 1) < p - 1$ for $n > 1$.*

For the proof of Theorem 1.1, we basically follow a beautiful argument of Abrashkin ([3]). We may assume $p \geq 3$ and $r \geq 1$. Consider the finite Galois extension

$$F_n = K(\pi^{1/p^n}, \zeta_{p^{n+1}})$$

of K whose upper ramification is bounded by $u(K, r, n)$. Let L_n be the finite Galois extension of F_n cut out by $T_{\text{st}, \bar{\pi}}^*(\mathcal{M})|_{G_{F_n}}$. Then we bound the ramification of L_n over K . For this, we show that to study this G_{F_n} -module we can use a variant over a smaller coefficient ring Σ of filtered (ϕ_r, N) -modules over S . In precise, we set

$$\Sigma = W[[u, E(u)^p/p]].$$

This ring Σ is small enough for the method of Abrashkin, in which he uses filtered modules of Fontaine-Laffaille ([13]) whose coefficient ring is W , to work also in the case where K is absolutely ramified.

Acknowledgments. The author would like to pay his gratitude to Iku Nakamura and Takeshi Saito for their warm encouragements. He wants to thank Manabu Yoshida for kindly allowing him to include the proof of Proposition 5.6. He also wants to thank Ahmed Abbès, Xavier Caruso, Takeshi Tsuji, and especially Victor Abrashkin and Yuichiro Taguchi, for useful discussions and comments. Finally, he is very grateful to the referee for his or her valuable comments to improve this paper.

2. FILTERED (ϕ_r, N) -MODULES OF BREUIL

In this section, we recall the theory of filtered (ϕ_r, N) -modules over S of Breuil, which is developed by himself and most recently by Caruso and Liu (see for example [6], [8], [17], [9]). In what follows, we always take the divided power envelope of a W -algebra with the compatibility condition with the natural divided power structure on pW .

Let p be a rational prime and σ be the Frobenius endomorphism of W . We fix once and for all a uniformizer π of K and a system $\{\pi_n\}_{n \in \mathbb{Z}_{\geq 0}}$ of p -power roots of π in \bar{K} such that $\pi_0 = \pi$ and $\pi_n = \pi_{n+1}^p$ for any n . Let $E(u)$

be the Eisenstein polynomial of π over W and set $S = (W[u]^{\text{PD}})^\wedge$, where PD means the divided power envelope and this is taken with respect to the ideal $(E(u))$, and \wedge means the p -adic completion. The ring S is endowed with the σ -semilinear endomorphism $\phi : u \mapsto u^p$ and a natural filtration $\text{Fil}^t S$ induced by the divided power structure such that $\phi(\text{Fil}^t S) \subseteq p^t S$ for $0 \leq t \leq p-1$. We set $\phi_t = p^{-t}\phi|_{\text{Fil}^t S}$ and $c = \phi_1(E(u)) \in S^\times$. Let N denote the W -linear derivation on S defined by the formula $N(u) = -u$. We also define a filtration, ϕ , ϕ_t and N on $S_n = S/p^n S$ similarly.

Let $r \in \{0, \dots, p-2\}$ be an integer. Set $'\text{Mod}_{/S}^{r,\phi,N}$ to be the category consisting of the following data:

- an S -module \mathcal{M} and its S -submodule $\text{Fil}^r \mathcal{M}$ containing $\text{Fil}^r S \cdot \mathcal{M}$,
- a ϕ -semilinear map $\phi_r : \text{Fil}^r \mathcal{M} \rightarrow \mathcal{M}$ satisfying

$$\phi_r(s_r m) = \phi_r(s_r) \phi(m)$$

- for any $s_r \in \text{Fil}^r S$ and $m \in \mathcal{M}$, where we set $\phi(m) = c^{-r} \phi_r(E(u)^r m)$,
- a W -linear map $N : \mathcal{M} \rightarrow \mathcal{M}$ such that
 - $N(sm) = N(s)m + sN(m)$ for any $s \in S$ and $m \in \mathcal{M}$,
 - $E(u)N(\text{Fil}^r \mathcal{M}) \subseteq \text{Fil}^r \mathcal{M}$,
 - the following diagram is commutative:

$$\begin{array}{ccc} \text{Fil}^r \mathcal{M} & \xrightarrow{\phi_r} & \mathcal{M} \\ E(u)N \downarrow & & \downarrow cN \\ \text{Fil}^r \mathcal{M} & \xrightarrow{\phi_r} & \mathcal{M}, \end{array}$$

and the morphisms of $'\text{Mod}_{/S}^{r,\phi,N}$ are defined to be the S -linear maps preserving Fil^r and commuting with ϕ_r and N . The category defined in the same way but dropping the data N is denoted by $'\text{Mod}_{/S}^{r,\phi}$. These categories have obvious notions of exact sequences. Let $\text{Mod}_{/S_1}^{r,\phi,N}$ denote the full subcategory of $'\text{Mod}_{/S}^{r,\phi,N}$ consisting of \mathcal{M} such that \mathcal{M} is free of finite rank over S_1 and generated as an S_1 -module by the image of ϕ_r . We write $\text{Mod}_{/S_\infty}^{r,\phi,N}$ for the smallest full subcategory which contains $\text{Mod}_{/S_1}^{r,\phi,N}$ and is stable under extensions. We let $\text{Mod}_{/S}^{r,\phi,N}$ denote the full subcategory consisting of \mathcal{M} such that

- the S -module \mathcal{M} is free of finite rank and generated by the image of ϕ_r ,
- the quotient $\mathcal{M}/\text{Fil}^r \mathcal{M}$ is p -torsion free.

We define full subcategories $\text{Mod}_{/S_1}^{r,\phi}$, $\text{Mod}_{/S_\infty}^{r,\phi}$ and $\text{Mod}_{/S}^{r,\phi}$ of $'\text{Mod}_{/S}^{r,\phi}$ in a similar way. For $\hat{\mathcal{M}} \in \text{Mod}_{/S}^{r,\phi,N}$ (resp. $\text{Mod}_{/S}^{r,\phi}$), the quotient $\hat{\mathcal{M}}/p^n \hat{\mathcal{M}}$ has a natural structure as an object of $\text{Mod}_{/S_\infty}^{r,\phi,N}$ (resp. $\text{Mod}_{/S_\infty}^{r,\phi}$).

For p -torsion objects, we also have the following categories. Consider the k -algebra $k[u]/(u^{ep}) \cong S_1/\text{Fil}^p S_1$ and let this algebra be denoted by \tilde{S}_1 . The algebra \tilde{S}_1 is equipped with the natural filtration, ϕ and N induced by those of S . Namely, $\text{Fil}^t \tilde{S}_1 = u^{et} \tilde{S}_1$, $\phi(u) = u^p$ and $N(u) = -u$. Let $'\text{Mod}_{/\tilde{S}_1}^{r,\phi,N}$ denote the category consisting of the following data:

- an \tilde{S}_1 -module $\tilde{\mathcal{M}}$ and its \tilde{S}_1 -submodule $\text{Fil}^r \tilde{\mathcal{M}}$ containing $u^{er} \tilde{\mathcal{M}}$,
- a ϕ -semilinear map $\phi_r : \text{Fil}^r \tilde{\mathcal{M}} \rightarrow \tilde{\mathcal{M}}$,
- a k -linear map $N : \tilde{\mathcal{M}} \rightarrow \tilde{\mathcal{M}}$ such that
 - $N(sm) = N(s)m + sN(m)$ for any $s \in \tilde{S}_1$ and $m \in \tilde{\mathcal{M}}$,
 - $u^e N(\text{Fil}^r \tilde{\mathcal{M}}) \subseteq \text{Fil}^r \tilde{\mathcal{M}}$,
 - the following diagram is commutative:

$$\begin{array}{ccc} \text{Fil}^r \tilde{\mathcal{M}} & \xrightarrow{\phi_r} & \tilde{\mathcal{M}} \\ u^e N \downarrow & & \downarrow cN \\ \text{Fil}^r \tilde{\mathcal{M}} & \xrightarrow{\phi_r} & \tilde{\mathcal{M}}, \end{array}$$

and whose morphisms are defined as before. Its full subcategory $\text{Mod}_{/\tilde{S}_1}^{r,\phi,N}$ is defined by the following condition:

- As an \tilde{S}_1 -module, $\tilde{\mathcal{M}}$ is free of finite rank and generated by the image of ϕ_r .

We define categories $'\text{Mod}_{/\tilde{S}_1}^{r,\phi}$ and $\text{Mod}_{/\tilde{S}_1}^{r,\phi}$ similarly. Then we can show as in the proof of [4, Proposition 2.2.2.1] that the natural functor $\mathcal{M} \mapsto \mathcal{M}/\text{Fil}^p S \cdot \mathcal{M}$ induces equivalences of categories $T : \text{Mod}_{/S_1}^{r,\phi,N} \rightarrow \text{Mod}_{/\tilde{S}_1}^{r,\phi,N}$ and $T_0 : \text{Mod}_{/S_1}^{r,\phi} \rightarrow \text{Mod}_{/\tilde{S}_1}^{r,\phi}$.

For $r = 0$, let $\text{Mod}_{/W_\infty}^\phi$ be the category consisting of the following data:

- a finite torsion W -module \tilde{M} ,
- a σ -semilinear automorphism $\phi : \tilde{M} \rightarrow \tilde{M}$.

Let κ be the kernel of the natural surjection $S \rightarrow W$ defined by $u \mapsto 0$. Since $\text{Tor}_1^S(\mathcal{M}, S/\kappa S) = 0$ for any $\mathcal{M} \in \text{Mod}_{/S_\infty}^{0,\phi}$, the proofs of [8, Lemme 2.2.7, Proposition 2.2.8] work also for the category $\text{Mod}_{/S_\infty}^{0,\phi,N}$ and we have a commutative diagram of categories

$$\begin{array}{ccc} \text{Mod}_{/S_\infty}^{0,\phi,N} & \longrightarrow & \text{Mod}_{/S_\infty}^{0,\phi} \\ & \searrow & \downarrow \\ & & \text{Mod}_{/W_\infty}^\phi, \end{array}$$

where the downward arrows and horizontal arrow are defined by $\mathcal{M} \mapsto \mathcal{M}/\kappa \mathcal{M}$ and forgetting N respectively and these three arrows are equivalences of categories.

Let A_{crys} and \hat{A}_{st} be p -adic period rings. These are constructed as follows. Put $\tilde{\mathcal{O}}_{\bar{K}} = \mathcal{O}_{\bar{K}}/p\mathcal{O}_{\bar{K}}$. Set R to be the ring

$$R = \varprojlim (\tilde{\mathcal{O}}_{\bar{K}} \leftarrow \tilde{\mathcal{O}}_{\bar{K}} \leftarrow \cdots),$$

where every arrow is the p -power map. For an element $x = (x_i)_{i \in \mathbb{Z}_{\geq 0}} \in R$ and an integer $n \geq 0$, we set

$$x^{(n)} = \lim_{m \rightarrow \infty} \hat{x}_{n+m}^{p^m} \in \mathcal{O}_{\mathbb{C}},$$

where \hat{x}_i is a lift of x_i in $\mathcal{O}_{\bar{K}}$ and $\mathcal{O}_{\mathbb{C}}$ is the p -adic completion of $\mathcal{O}_{\bar{K}}$. Let v_p denote the valuation of $\mathcal{O}_{\mathbb{C}}$ normalized as $v_p(p) = 1$. Then the ring R is a complete valuation ring whose valuation of an element $x \in R$ is given by $v_R(x) = v_p(x^{(0)})$. We define a natural ring homomorphism θ by

$$\begin{aligned} \theta : W(R) &\rightarrow \mathcal{O}_{\mathbb{C}} \\ (x_0, x_1, \dots) &\mapsto \sum_{n \geq 0} p^n x_n^{(n)}. \end{aligned}$$

Then A_{crys} is the p -adic completion of the divided power envelope of $W(R)$ with respect to the principal ideal $\text{Ker}(\theta)$ and \hat{A}_{st} is the p -adic completion of the divided power polynomial ring $A_{\text{crys}}\langle X \rangle$ over A_{crys} . We set $A_{\text{crys},\infty} = A_{\text{crys}} \otimes_W K_0/W$ and $\hat{A}_{\text{st},\infty} = \hat{A}_{\text{st}} \otimes_W K_0/W$. Put $\underline{\pi} = (\pi_n)_{n \in \mathbb{Z}_{\geq 0}} \in R$, where we abusively let π_n denote the image of $\pi_n \in \mathcal{O}_{\bar{K}}$ in $\tilde{\mathcal{O}}_{\bar{K}}$. These rings are considered as S -algebras by the ring homomorphisms $S \rightarrow \hat{A}_{\text{st}}$ and $\hat{A}_{\text{st}} \rightarrow A_{\text{crys}}$ which are defined by $u \mapsto [\underline{\pi}]/(1+X)$ and $X \mapsto 0$, respectively. The ring A_{crys} is endowed with a natural filtration induced by the divided power structure, a natural Frobenius endomorphism ϕ and the ϕ -semilinear map $\phi_t = p^{-t}\phi|_{\text{Fil}^t A_{\text{crys}}}$. With these structures, A_{crys} and $A_{\text{crys},\infty}$ are considered as objects of $'\text{Mod}_{/S}^{r,\phi}$. Moreover, the absolute Galois group G_K acts naturally on these two rings. As for \hat{A}_{st} , its filtration is defined by

$$\text{Fil}^t \hat{A}_{\text{st}} = \left\{ \sum_{i \geq 0} a_i \frac{X^i}{i!} \mid a_i \in \text{Fil}^{t-i} A_{\text{crys}}, \lim_{i \rightarrow \infty} a_i = 0 \right\}$$

and the Frobenius structure of A_{crys} extends to \hat{A}_{st} by

$$\begin{aligned} \phi(X) &= (1+X)^p - 1, \\ \phi_t &= p^{-t}\phi|_{\text{Fil}^t \hat{A}_{\text{st}}}. \end{aligned}$$

We write N also for the A_{crys} -linear derivation on \hat{A}_{st} defined by $N(X) = 1+X$. The rings \hat{A}_{st} and $\hat{A}_{\text{st},\infty}$ are objects of $'\text{Mod}_{/S}^{r,\phi,N}$. The G_K -action on A_{crys} naturally extends to an action on \hat{A}_{st} . Indeed, the action of $g \in G_K$ on \hat{A}_{st} is defined by the formula

$$g(X) = [\underline{\varepsilon}(g)](1+X) - 1,$$

where $g(\pi_n) = \varepsilon_n(g)\pi_n$ and $\underline{\varepsilon}(g) = (\varepsilon_n(g))_{n \in \mathbb{Z}_{\geq 0}} \in R$ with the abusive notation as above.

These rings have other descriptions, as follows. For an integer $n \geq 1$, put $W_n = W/p^n W$ and let $W_n(\tilde{\mathcal{O}}_{\bar{K}})$ be the ring of Witt vectors of length n associated to $\tilde{\mathcal{O}}_{\bar{K}}$. We define a W_n -algebra structure on $W_n(\tilde{\mathcal{O}}_{\bar{K}})$ by twisting the natural W_n -algebra structure by σ^{-n} . Then the natural ring homomorphism

$$\begin{aligned} \theta_n : W_n(\tilde{\mathcal{O}}_{\bar{K}}) &\rightarrow \mathcal{O}_{\bar{K}}/p^n \mathcal{O}_{\bar{K}} \\ (a_0, \dots, a_{n-1}) &\mapsto \sum_{i=0}^{n-1} p^i \hat{a}_i^{p^{n-i}}, \end{aligned}$$

where \hat{a}_i is a lift of a_i in $\mathcal{O}_{\bar{K}}$, is W_n -linear. Let us denote $W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}})$ the divided power envelope of $W_n(\tilde{\mathcal{O}}_{\bar{K}})$ with respect to the ideal $\text{Ker}(\theta_n)$. This ring is considered as an S -algebra by $u \mapsto [\pi_n]$. This ring also has a natural filtration defined by the divided power structure, and a natural G_K -module structure. The Frobenius endomorphism of the ring of Witt vectors induces on this ring a ϕ -semilinear Frobenius endomorphism, which is denoted also by ϕ . Then, by the S -linear transition maps

$$\begin{aligned} W_{n+1}^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}}) &\rightarrow W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}}) \\ (a_0, \dots, a_n) &\mapsto (a_0^p, \dots, a_{n-1}^p), \end{aligned}$$

these S -algebras form a projective system compatible with all the structures. Using this transition map, a ϕ -semilinear map

$$\phi_r : \text{Fil}^r W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}}) \rightarrow W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}})$$

is defined by setting $\phi_r(x)$ to be the image of $p^{-r}\phi(\hat{x})$, where \hat{x} is a lift of x in $\text{Fil}^r W_{n+r}^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}})$. By definition, the maps ϕ_r are also compatible with the transition maps. The S -algebra $W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}})$ is considered as an object of $'\text{Mod}_{/S}^{r,\phi}$. Then we have a natural isomorphism in $'\text{Mod}_{/S}^{r,\phi}$

$$\begin{aligned} A_{\text{crys}}/p^n A_{\text{crys}} &\rightarrow W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}}) \\ (x_0, \dots, x_{n-1}) &\mapsto (x_{0,n}, \dots, x_{n-1,n}), \end{aligned}$$

where we set $x_i = (x_{i,k})_{k \in \mathbb{Z}_{\geq 0}}$ for $x_i \in R$.

Similarly, the divided power polynomial ring $W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}})\langle X \rangle$ over $W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}})$ is considered as an S -algebra by $u \mapsto [\pi_n]/(1+X)$. This ring has a natural filtration coming from the divided power structure. We define a G_K -action on this ring by

$$g(X) = [\varepsilon_n(g)](1+X) - 1.$$

We also define a ϕ -semilinear Frobenius endomorphism, which we also write as ϕ , by $\phi(X) = (1+X)^p - 1$ and a $W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}})$ -linear derivation N by $N(X) = 1+X$. These rings form a projective system of S -algebras compatible with

all the structures by the transition maps defined by the maps above and $X \mapsto X$. We define ϕ -semilinear maps

$$\phi_r : \mathrm{Fil}^r W_n^{\mathrm{PD}}(\tilde{\mathcal{O}}_{\bar{K}})\langle X \rangle \rightarrow W_n^{\mathrm{PD}}(\tilde{\mathcal{O}}_{\bar{K}})\langle X \rangle$$

compatible with the transition maps as before. The S -algebra $W_n^{\mathrm{PD}}(\tilde{\mathcal{O}}_{\bar{K}})\langle X \rangle$ is considered as an object of $'\mathrm{Mod}_{/S}^{r,\phi,N}$ and there exists a natural isomorphism in $'\mathrm{Mod}_{/S}^{r,\phi,N}$

$$\begin{aligned} \hat{A}_{\mathrm{st}}/p^n \hat{A}_{\mathrm{st}} &\rightarrow W_n^{\mathrm{PD}}(\tilde{\mathcal{O}}_{\bar{K}})\langle X \rangle \\ (x_0, \dots, x_{n-1}) &\mapsto (x_{0,n}, \dots, x_{n-1,n}) \\ X &\mapsto X \end{aligned}$$

which is G_K -linear.

Put $K_n = K(\pi_n)$ and $K_\infty = \cup_n K_n$. For $\mathcal{M} \in \mathrm{Mod}_{/S_\infty}^{r,\phi,N}$, we define a G_K -module $T_{\mathrm{st},\underline{\pi}}^*(\mathcal{M})$ to be

$$T_{\mathrm{st},\underline{\pi}}^*(\mathcal{M}) = \mathrm{Hom}_{S, \mathrm{Fil}^r, \phi_r, N}(\mathcal{M}, \hat{A}_{\mathrm{st},\infty}).$$

When \mathcal{M} is killed by p^n , we have a natural identification of G_K -modules

$$T_{\mathrm{st},\underline{\pi}}^*(\mathcal{M}) = \mathrm{Hom}_{S, \mathrm{Fil}^r, \phi_r, N}(\mathcal{M}, W_n^{\mathrm{PD}}(\tilde{\mathcal{O}}_{\bar{K}})\langle X \rangle).$$

Note that the G_K -module on the right-hand side is independent of the choice of π_k for $k > n$. Since the natural map

$$\begin{aligned} W_n^{\mathrm{PD}}(\tilde{\mathcal{O}}_{\bar{K}})\langle X \rangle &\rightarrow W_n^{\mathrm{PD}}(\tilde{\mathcal{O}}_{\bar{K}}) \\ X &\mapsto 0 \end{aligned}$$

is G_{K_n} -linear, we also have a G_{K_n} -linear isomorphism ([6, Lemme 2.3.1.1])

$$T_{\mathrm{st},\underline{\pi}}^*(\mathcal{M})|_{G_{K_n}} \rightarrow \mathrm{Hom}_{S, \mathrm{Fil}^r, \phi_r}(\mathcal{M}, W_n^{\mathrm{PD}}(\tilde{\mathcal{O}}_{\bar{K}})).$$

On the other hand, for $r = 0$, the proof of [8, Proposition 2.3.13] shows that the G_K -module $T_{\mathrm{st},\underline{\pi}}^*(\mathcal{M})$ is unramified for any $\mathcal{M} \in \mathrm{Mod}_{/S_\infty}^{0,\phi,N}$.

A variant of filtered (ϕ_r, N) -modules over S is also introduced by Breuil and Kisin, and developed also by Caruso and Liu (see for example [15], [16], [17], [9]). Put $\mathfrak{S} = W[[u]]$ and let $\phi : \mathfrak{S} \rightarrow \mathfrak{S}$ be the σ -semilinear Frobenius endomorphism defined by $\phi(u) = u^p$. Let $'\mathrm{Mod}_{/\mathfrak{S}}^{r,\phi}$ denote the category consisting of the following data:

- an \mathfrak{S} -module \mathfrak{M} ,
- a ϕ -semilinear map $\mathfrak{M} \rightarrow \mathfrak{M}$, which is denoted also by ϕ , such that the cokernel of the map $1 \otimes \phi : \phi^* \mathfrak{M} \rightarrow \mathfrak{M}$, where we set $\phi^* \mathfrak{M} = \mathfrak{S} \otimes_{\phi, \mathfrak{S}} \mathfrak{M}$, is killed by $E(u)^r$,

and whose morphisms are defined as before. The full subcategory of $'\mathrm{Mod}_{/\mathfrak{S}}^{r,\phi}$ consisting of \mathfrak{M} such that \mathfrak{M} is free of finite rank over $\mathfrak{S}/p\mathfrak{S}$ (*resp.* over \mathfrak{S}) is denoted by $\mathrm{Mod}_{/\mathfrak{S}_1}^{r,\phi}$ (*resp.* $\mathrm{Mod}_{/\mathfrak{S}}^{r,\phi}$). We let $\mathrm{Mod}_{/\mathfrak{S}_\infty}^{r,\phi}$ denote the smallest full subcategory which contains $\mathrm{Mod}_{/\mathfrak{S}_1}^{r,\phi}$ and is stable under extensions, as

before. Then we have an exact functor ([9, Proposition 2.1.2], see also [15, Proposition 1.1.11])

$$\mathcal{M}_{\mathfrak{S}_\infty} : \text{Mod}_{/\mathfrak{S}_\infty}^{r,\phi} \rightarrow \text{Mod}_{/S_\infty}^{r,\phi}.$$

For $\mathfrak{M} \in \text{Mod}_{/\mathfrak{S}_\infty}^{r,\phi}$, the filtered ϕ_r -module $\mathcal{M} = \mathcal{M}_{\mathfrak{S}_\infty}(\mathfrak{M})$ over S is defined as follows:

- $\mathcal{M} = S \otimes_{\phi,\mathfrak{S}} \mathfrak{M}$,
- $\text{Fil}^r \mathcal{M} = \text{Ker}(\mathcal{M} \xrightarrow{1 \otimes \phi} S \otimes_{\mathfrak{S}} \mathfrak{M} \rightarrow (S/\text{Fil}^r S) \otimes_{\mathfrak{S}} \mathfrak{M})$,
- $\phi_r : \text{Fil}^r \mathcal{M} \xrightarrow{1 \otimes \phi} \text{Fil}^r S \otimes_{\mathfrak{S}} \mathfrak{M} \xrightarrow{\phi_r \otimes 1} S \otimes_{\phi,\mathfrak{S}} \mathfrak{M} = \mathcal{M}$.

We write $\mathcal{M}_{\mathfrak{S}}$ for the functor $\text{Mod}_{/\mathfrak{S}}^{r,\phi} \rightarrow \text{Mod}_{/S}^{r,\phi}$ defined similarly.

3. FILTERED ϕ_r -MODULES OVER Σ

In this section, we define another variant $\text{Mod}_{/\Sigma_\infty}^{r,\phi}$ of the category $\text{Mod}_{/S_\infty}^{r,\phi}$ over a subring Σ of the ring S , and prove that they are categorically equivalent.

Let p be a rational prime and r be an integer such that $0 \leq r < p-1$. Consider the W -algebra $\Sigma = W[[u, Y]]/(E(u)^p - pY)$ as in [6, Subsection 3.2]. We regard Σ as a subring of S by the map sending Y to $E(u)^p/p$. Then the element $c = \phi_1(E(u)) \in S^\times$ is contained in Σ^\times . We define on Σ a σ -semilinear Frobenius endomorphism ϕ by $\phi(u) = u^p$ and $\phi(Y) = p^{p-1}c^p$. Put $\text{Fil}^t \Sigma = (E(u)^t, Y)$ for $0 \leq t \leq p-1$ and $\text{Fil}^p \Sigma = (Y)$. Then we have $\phi(\text{Fil}^t \Sigma) \subseteq p^t \Sigma$ for $0 \leq t \leq p-1$. We put $\phi_t = p^{-t} \phi|_{\text{Fil}^t \Sigma}$. We also set $\Sigma_n = \Sigma/p^n \Sigma$ and put on this ring the natural structures induced by those of Σ .

We define a category $'\text{Mod}_{/\Sigma}^{r,\phi}$ of filtered ϕ_r -modules over Σ to be the category consisting of the following data:

- a Σ -module M and its Σ -submodule $\text{Fil}^r M$ containing $\text{Fil}^r \Sigma \cdot M$,
- a ϕ -semilinear map $\phi_r : \text{Fil}^r M \rightarrow M$ satisfying $\phi_r(s_r m) = \phi_r(s_r) \phi(m)$ for any $s_r \in \text{Fil}^r \Sigma$ and $m \in M$, where we set $\phi(m) = c^{-r} \phi_r(E(u)^r m)$,

and whose morphisms are defined in the same manner as $'\text{Mod}_{/S}^{r,\phi}$. This category has a natural notion of exact sequences. We define its full subcategory $\text{Mod}_{/\Sigma_1}^{r,\phi}$ to be the category consisting of M which is free of finite rank and generated by the image of ϕ_r as a Σ_1 -module. We also let $\text{Mod}_{/\Sigma_\infty}^{r,\phi}$ denote the smallest full subcategory of $'\text{Mod}_{/\Sigma}^{r,\phi}$ which contains $\text{Mod}_{/\Sigma_1}^{r,\phi}$ and is stable under extensions. Moreover, we define a full subcategory $\text{Mod}_{/\Sigma}^{r,\phi}$ of $'\text{Mod}_{/\Sigma}^{r,\phi}$ to be the category consisting of M such that

- the Σ -module M is free of finite rank and generated by the image of ϕ_r ,
- the quotient $M/\text{Fil}^r M$ is p -torsion free.

Then we see that for $\hat{M} \in \text{Mod}_{/\Sigma}^{r,\phi}$, the quotient $\hat{M}/p^n \hat{M}$ is naturally considered as an object of $\text{Mod}_{/\Sigma_\infty}^{r,\phi}$.

The natural ring isomorphism $\Sigma_1/\text{Fil}^p \Sigma_1 \cong \tilde{S}_1$ defines a functor $T_{0,\Sigma} : \text{Mod}_{/\Sigma_1}^{r,\phi} \rightarrow \text{Mod}_{/\tilde{S}_1}^{r,\phi}$ by $M \mapsto M/\text{Fil}^p \Sigma_1 \cdot M$. Then just as in the case of the functor $T_0 : \text{Mod}_{/S_1}^{r,\phi} \rightarrow \text{Mod}_{/\tilde{S}_1}^{r,\phi}$ ([4, Proposition 2.2.2.1]), we can show the following lemma.

Lemma 3.1. *The functor $T_{0,\Sigma} : \text{Mod}_{/\Sigma_1}^{r,\phi} \rightarrow \text{Mod}_{/\tilde{S}_1}^{r,\phi}$ is an equivalence of categories.*

On the other hand, [6, Proposition 2.2.1.3] and Nakayama's lemma show the following.

Lemma 3.2. *Let M be an object of $\text{Mod}_{/\Sigma_1}^{r,\phi}$ of rank d over Σ_1 . Then there exists a basis $\{e_1, \dots, e_d\}$ of M such that $\text{Fil}^r M = \Sigma_1 u^{r_1} e_1 \oplus \dots \oplus \Sigma_1 u^{r_d} e_d + \text{Fil}^p \Sigma_1 \cdot M$ for some integers r_1, \dots, r_d with $0 \leq r_i \leq er$ for any i .*

Then we can show the following lemma just as in the proof of [6, Lemme 2.3.1.3].

Lemma 3.3. *The functor*

$$M \mapsto \text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, A_{\text{crys}, \infty})$$

from $\text{Mod}_{/\Sigma_\infty}^{r,\phi}$ to the category of G_{K_∞} -modules is exact.

For $M \in \text{Mod}_{/\Sigma_1}^{r,\phi}$, we can show as in the case of the category $\text{Mod}_{/S_1}^{r,\phi}$ that there is an isomorphism of G_{K_1} -modules

$$\text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, (\tilde{\mathcal{O}}_{\bar{K}})^{\text{PD}}) \rightarrow \text{Hom}_{\tilde{S}_1, \text{Fil}^r, \phi_r}(T_{0,\Sigma}(M), \tilde{\mathcal{O}}_{\bar{K}}),$$

where $\tilde{\mathcal{O}}_{\bar{K}}$ is considered as an object of $\text{Mod}_{/\tilde{S}_1}^{r,\phi}$ by the natural isomorphism

$$(\tilde{\mathcal{O}}_{\bar{K}})^{\text{PD}}/\text{Fil}^p(\tilde{\mathcal{O}}_{\bar{K}})^{\text{PD}} \rightarrow \tilde{\mathcal{O}}_{\bar{K}}.$$

Thus [6, Lemme 2.3.1.2] implies the following.

Lemma 3.4. *For $M \in \text{Mod}_{/\Sigma_1}^{r,\phi}$, we have*

$$\#\text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, (\tilde{\mathcal{O}}_{\bar{K}})^{\text{PD}}) = p^d,$$

where $d = \dim_{\Sigma_1} M$.

For the category $\text{Mod}_{/\Sigma_\infty}^{r,\phi}$, we have the following lemma.

Lemma 3.5. *Let M be in $\text{Mod}_{/\Sigma_\infty}^{r,\phi}$. Then there exists $\alpha_1, \dots, \alpha_d \in \text{Fil}^r M$ such that $\text{Fil}^r M = \Sigma \alpha_1 + \dots + \Sigma \alpha_d + \text{Fil}^p \Sigma \cdot M$ and the elements $e_1 = \phi_r(\alpha_1), \dots, e_d = \phi_r(\alpha_d)$ form a system of generators of M .*

Proof. By induction and Lemma 3.2, we may assume that there exists an exact sequence of the category $\text{Mod}_{/\Sigma_\infty}^{r,\phi}$

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

such that the lemma holds for M' and M'' . Let $\alpha'_1, \dots, \alpha'_{l'}$ (*resp.* $\alpha''_1, \dots, \alpha''_{l''}$) be elements of $\text{Fil}^r M'$ (*resp.* $\text{Fil}^r M''$) as in the lemma. Let $\alpha_l \in \text{Fil}^r M$ be a lift of α'_l . Then the elements $\alpha'_1, \dots, \alpha'_{l'}, \alpha_1, \dots, \alpha_{l''}$ satisfy the condition in the lemma for M . \square

Corollary 3.6. *Let M be an object of $\text{Mod}_{/\Sigma_\infty}^{r,\phi}$ and $C \in M_d(\Sigma)$ be a matrix satisfying*

$$(\alpha_1, \dots, \alpha_d) = (e_1, \dots, e_d)C$$

with the notation of the previous lemma. Let A be an object of $'\text{Mod}_{/\Sigma}^{r,\phi}$. Then a Σ -linear homomorphism $f : M \rightarrow A$ preserving Fil^r also commutes with ϕ_r if and only if

$$\phi_r(f(e_1, \dots, e_d)C) = (f(e_1), \dots, f(e_d)).$$

Proof. Suppose that the latter condition holds. Then we have $\phi_r(f(\alpha_i)) = f(\phi_r(\alpha_i))$ for any i . We only have to check the equality $\phi_r \circ f = f \circ \phi_r$ on $\text{Fil}^p \Sigma \cdot M$. Suppose that this equality holds on the submodule $p^{l+1} \text{Fil}^p \Sigma \cdot M$. For $m \in M$, we can take $m' \in \text{Fil}^p \Sigma \cdot M$ such that $E(u)^r m = \sum_i s_i \alpha_i + m'$. Let s be in $\text{Fil}^p \Sigma$. Then we have

$$f(\phi_r(p^l s m)) = p^l \phi_r(s) c^{-r} \sum_i \phi(s_i) f(\phi_r(\alpha_i)) + p^l \phi_r(s) c^{-r} f(\phi_r(m')).$$

Since $\phi_r(\text{Fil}^p \Sigma) \subseteq p\Sigma$, this equals to $\phi_r(f(p^l s m))$ by assumption. Thus the lemma follows by induction. \square

Corollary 3.7. *Let M and A be as above and $J \subseteq \text{Fil}^r A$ be a Σ -submodule of A such that $\phi_r(J) \subseteq J$. We can consider the Σ -module A/J naturally as an object of $'\text{Mod}_{/\Sigma}^{r,\phi}$. Suppose that for any $x \in J$, there exists $t \in \mathbb{Z}_{\geq 0}$ such that $\phi_r^t(x) = 0$. Then the natural homomorphism of abelian groups*

$$\text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, A) \rightarrow \text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, A/J)$$

is an isomorphism.

Proof. The proof is similar to [3, Subsection 2.2]. We consider the Σ -submodule J as an object of the category $'\text{Mod}_{/\Sigma}^{r,\phi}$ by putting $\text{Fil}^r J = J$. By devissage, it is enough to show that, for any $M \in \text{Mod}_{/\Sigma_1}^{r,\phi}$, we have $\text{Ext}_{\text{Mod}_{/\Sigma}^{r,\phi}}(M, J) = 0$ and the map in the corollary is an isomorphism. For the first assertion, let

$$0 \longrightarrow J \longrightarrow \mathcal{E} \longrightarrow M \longrightarrow 0$$

be an extension in the category $'\text{Mod}_{/\Sigma}^{r,\phi}$. Let e_i, α_i and C be as in Corollary 3.6 such that e_1, \dots, e_d form a basis of M . Let $\hat{e}_i \in \mathcal{E}$ be a lift of $e_i \in M$. Then we have $(\hat{e}_1, \dots, \hat{e}_d)C \in (\text{Fil}^r \mathcal{E})^{\oplus d}$ and

$$\phi_r((\hat{e}_1, \dots, \hat{e}_d)C) = (\hat{e}_1 + \delta_1, \dots, \hat{e}_d + \delta_d)$$

for some $\delta_1, \dots, \delta_d \in J$. On the other hand, there exists a unique d -tuple $(x_1, \dots, x_d) \in J^{\oplus d}$ satisfying the equation

$$\phi_r((\hat{e}_1 + x_1, \dots, \hat{e}_d + x_d)C) = (\hat{e}_1 + x_1, \dots, \hat{e}_d + x_d).$$

Indeed, the d -tuple

$$\sum_{i=0}^t (\phi_r^i(\delta_1), \dots, \phi_r^i(\delta_d)) \phi(C) \cdots \phi^{i-1}(C) \phi^i(C)$$

is stable for sufficiently large t by assumption and this limit gives a unique solution of the equation. Then we have

$$(p(\hat{e}_1 + x_1), \dots, p(\hat{e}_d + x_d)) = \phi_r(p(\hat{e}_1 + x_1), \dots, p(\hat{e}_d + x_d)) \phi(C).$$

Since the d -tuple on the left-hand side is contained in $J^{\oplus d}$, we see that this d -tuple is zero and $e_i \mapsto \hat{e}_i + x_i$ defines a section $M \rightarrow \mathcal{E}$. We can prove the second assertion similarly. \square

Next we show that the two categories $\text{Mod}_{/\Sigma_\infty}^{r,\phi}$ and $\text{Mod}_{/S_\infty}^{r,\phi}$ are in fact equivalent. For $M \in \text{Mod}_{/\Sigma_\infty}^{r,\phi}$, we associate to it an S -module \mathcal{M} by setting $\mathcal{M} = S \otimes_\Sigma M$. We also define its S -submodule $\text{Fil}^r \mathcal{M}$ by

$$\text{Fil}^r \mathcal{M} = \text{Ker}(\mathcal{M} = S \otimes_\Sigma M \rightarrow S/\text{Fil}^r S \otimes_\Sigma M/\text{Fil}^r M \simeq M/\text{Fil}^r M),$$

where the last isomorphism is induced by the natural isomorphisms of W -algebras

$$W[u]/(E(u)^r) \rightarrow \Sigma/\text{Fil}^r \Sigma \rightarrow S/\text{Fil}^r S.$$

These associations induce two functors from $\text{Mod}_{/\Sigma_\infty}^{r,\phi}$ to the category of S -modules, $M \mapsto \mathcal{M}$ and $M \mapsto \text{Fil}^r \mathcal{M}$. Since the rings S and $W[u]/(E(u)^r)$ are p -torsion free, we have $\text{Tor}_1^\Sigma(\Sigma_1, S) = \text{Tor}_1^\Sigma(\Sigma_1, \Sigma/\text{Fil}^r \Sigma) = 0$ and thus $\text{Tor}_1^\Sigma(M, S) = \text{Tor}_1^\Sigma(M, \Sigma/\text{Fil}^r \Sigma) = 0$ for any $M \in \text{Mod}_{/\Sigma_\infty}^{r,\phi}$. Hence we see that these two functors are exact.

We define $\phi_r : \text{Fil}^r \mathcal{M} \rightarrow \mathcal{M}$ as follows. Note that $\text{Fil}^r S \otimes_\Sigma M \subseteq \mathcal{M}$ and $\text{Fil}^r \mathcal{M}$ is equal to $\text{Fil}^r S \otimes_\Sigma M + \text{Im}(S \otimes_\Sigma \text{Fil}^r M \rightarrow \mathcal{M})$. Set $\phi_r' : \text{Fil}^r S \otimes_\Sigma M \rightarrow \mathcal{M}$ to be $\phi_r' = \phi_r \otimes \phi$.

Lemma 3.8. *The map $\phi \otimes \phi_r : S \otimes_\Sigma \text{Fil}^r M \rightarrow \mathcal{M}$ induces a ϕ -semilinear map $\phi_r'' : \text{Im}(S \otimes_\Sigma \text{Fil}^r M \rightarrow \mathcal{M}) \rightarrow \mathcal{M}$.*

Proof. Let $z = \sum_i s_i \otimes m_i$ be in $S \otimes_\Sigma \text{Fil}^r M$ with $s_i \in S$ and $m_i \in \text{Fil}^r M$. Let \bar{z} be its image in \mathcal{M} and suppose that $\bar{z} = 0$. Write $s_i = s_i' + s_i''$ with $s_i' \in \Sigma$ and $s_i'' \in \text{Fil}^p S$. Since we have an isomorphism $\mathcal{M}/\text{Fil}^r S \cdot \mathcal{M} \simeq M/\text{Fil}^r \Sigma \cdot M$,

we can find elements $s^{(j)} \in \text{Fil}^r \Sigma$ and $m^{(j)} \in M$ such that the equality $\sum_i s'_i m_i = \sum_j s^{(j)} m^{(j)}$ holds in M . Then we have

$$0 = \bar{z} = \sum_i 1 \otimes s'_i m_i + \sum_i s''_i \otimes m_i = \sum_j s^{(j)} \otimes m^{(j)} + \sum_i s''_i \otimes m_i$$

in \mathcal{M} . On the other hand, the element $(\phi \otimes \phi_r)(z) \in \mathcal{M}$ is equal to

$$\sum_j 1 \otimes \phi_r(s^{(j)} m^{(j)}) + \sum_i \phi(s''_i) \otimes \phi_r(m_i).$$

Since $\phi = p^r \phi_r$, this equals $\phi'_r(\sum_j s^{(j)} \otimes m^{(j)} + \sum_i s''_i \otimes m_i) = 0$. \square

Lemma 3.9. *The maps ϕ'_r and ϕ''_r patch together and define a ϕ -semilinear map $\phi_r : \text{Fil}^r \mathcal{M} \rightarrow \mathcal{M}$.*

Proof. Since ϕ'_r and ϕ''_r coincide on $\text{Im}(\text{Fil}^r S \otimes_\Sigma \text{Fil}^r M \rightarrow \mathcal{M})$, it is enough to show that $1 \otimes \phi_r(m) = \phi'_r(\sum_i s_i \otimes m_i)$ for any $m \in \text{Fil}^r M$, $s_i \in \text{Fil}^r S$ and $m_i \in M$ satisfying $1 \otimes m = \sum_i s_i \otimes m_i$ in \mathcal{M} . As in the proof of Lemma 3.8, the element m can be written as $m = \sum_j s^{(j)} m^{(j)}$ for some $s^{(j)} \in \text{Fil}^r \Sigma$ and $m^{(j)} \in M$. By assumption, we have $\sum_i s_i \otimes m_i = \sum_j s^{(j)} \otimes m^{(j)}$ in $\text{Fil}^r S \otimes_\Sigma M$. Hence the lemma follows. \square

Then we see that this construction defines a functor $\mathcal{M}_{\Sigma_\infty} : \text{Mod}_{/\Sigma_\infty}^{r,\phi} \rightarrow \text{Mod}_{/S_\infty}^{r,\phi}$.

Lemma 3.10. *The functor $\mathcal{M}_{\Sigma_\infty}$ induces an equivalence of categories $\text{Mod}_{/\Sigma_1}^{r,\phi} \rightarrow \text{Mod}_{/S_1}^{r,\phi}$.*

Proof. Consider the diagram of functors

$$\begin{array}{ccc} \text{Mod}_{/\Sigma_1}^{r,\phi} & \xrightarrow{\mathcal{M}_{\Sigma_\infty}} & \text{Mod}_{/S_1}^{r,\phi} \\ & \searrow T_{0,\Sigma} & \downarrow T_0 \\ & & \text{Mod}_{/\tilde{S}_1}^{r,\phi}. \end{array}$$

From the definition, we see that this diagram is commutative. By Lemma 3.1, the downward arrows are equivalences of categories. Thus the lemma follows. \square

Then a devissage argument as in [15, Proposition 1.1.11] shows the following corollary.

Corollary 3.11. *The functor $\mathcal{M}_{\Sigma_\infty} : \text{Mod}_{/\Sigma_\infty}^{r,\phi} \rightarrow \text{Mod}_{/S_\infty}^{r,\phi}$ is fully faithful.*

To show the essential surjectivity of the functor $\mathcal{M}_{\Sigma_\infty}$, we define another functor $M_{\mathfrak{S}_\infty} : \text{Mod}_{/\mathfrak{S}_\infty}^{r,\phi} \rightarrow \text{Mod}_{/\Sigma_\infty}^{r,\phi}$ which is defined in a similar way to the functor $\mathcal{M}_{\mathfrak{S}_\infty} : \text{Mod}_{/\mathfrak{S}_\infty}^{r,\phi} \rightarrow \text{Mod}_{/S_\infty}^{r,\phi}$. For an \mathfrak{S} -module \mathfrak{M} in $\text{Mod}_{/\mathfrak{S}_\infty}^{r,\phi}$, we associate to it a Σ -module $M \in \text{Mod}_{/\Sigma}^{r,\phi}$ as follows:

- $M = \Sigma \otimes_{\phi, \mathfrak{S}} \mathfrak{M}$,
- $\text{Fil}^r M = \text{Ker}(M \xrightarrow{1 \otimes \phi} \Sigma \otimes_{\mathfrak{S}} \mathfrak{M} \rightarrow (\Sigma/\text{Fil}^r \Sigma) \otimes_{\mathfrak{S}} \mathfrak{M})$,
- $\phi_r : \text{Fil}^r M \xrightarrow{1 \otimes \phi} \text{Fil}^r \Sigma \otimes_{\mathfrak{S}} \mathfrak{M} \xrightarrow{\phi_r \otimes 1} \Sigma \otimes_{\phi, \mathfrak{S}} \mathfrak{M} = M$.

We can check that this defines an exact functor $\text{Mod}_{/\mathfrak{S}_{\infty}}^{r, \phi} \rightarrow \text{Mod}_{/\Sigma_{\infty}}^{r, \phi}$ as in the proof of [15, Proposition 1.1.11]. We let this functor be denoted by $M_{\mathfrak{S}_{\infty}}$.

Lemma 3.12. *The diagram of functors*

$$\begin{array}{ccc} \text{Mod}_{/\mathfrak{S}_{\infty}}^{r, \phi} & \xrightarrow{M_{\mathfrak{S}_{\infty}}} & \text{Mod}_{/\Sigma_{\infty}}^{r, \phi} \\ & \searrow \mathcal{M}_{\mathfrak{S}_{\infty}} & \downarrow \mathcal{M}_{\Sigma_{\infty}} \\ & & \text{Mod}_{/S_{\infty}}^{r, \phi} \end{array}$$

is commutative.

Proof. For $\mathfrak{M} \in \text{Mod}_{/\mathfrak{S}_{\infty}}^{r, \phi}$, put $M = M_{\mathfrak{S}_{\infty}}(\mathfrak{M})$ and $\mathcal{M} = \mathcal{M}_{\mathfrak{S}_{\infty}}(\mathfrak{M})$. Then $\mathcal{M} = S \otimes_{\Sigma} M$ as an S -module. Let $\text{Fil}^r \mathcal{M}$ and $\phi_r : \text{Fil}^r \mathcal{M} \rightarrow \mathcal{M}$ denote the filtration and Frobenius structure defined by the functor $\mathcal{M}_{\mathfrak{S}_{\infty}}$. We also let $\hat{\text{Fil}}^r \mathcal{M}$ and $\hat{\phi}_r : \hat{\text{Fil}}^r \mathcal{M} \rightarrow \mathcal{M}$ denote those defined by $\mathcal{M}_{\Sigma_{\infty}}$.

The S -module $\text{Fil}^r \mathcal{M}$ contains $\hat{\text{Fil}}^r \mathcal{M}$. Conversely, let z be an element of $\text{Fil}^r \mathcal{M}$. Note that $\text{Fil}^p S \cdot \mathcal{M} \subseteq \hat{\text{Fil}}^r \mathcal{M}$. Thus, to show $z \in \hat{\text{Fil}}^r \mathcal{M}$, we may assume that $z \in \text{Im}(M \rightarrow \mathcal{M})$. Then the commutative diagram whose right vertical arrow is an isomorphism

$$\begin{array}{ccccc} M = \Sigma \otimes_{\phi, \mathfrak{S}} \mathfrak{M} & \xrightarrow{1 \otimes \phi} & \Sigma \otimes_{\mathfrak{S}} \mathfrak{M} & \longrightarrow & \Sigma/\text{Fil}^r \Sigma \otimes_{\mathfrak{S}} \mathfrak{M} \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{M} = S \otimes_{\phi, \mathfrak{S}} \mathfrak{M} & \xrightarrow{1 \otimes \phi} & S \otimes_{\mathfrak{S}} \mathfrak{M} & \longrightarrow & S/\text{Fil}^r S \otimes_{\mathfrak{S}} \mathfrak{M} \end{array}$$

implies that $z \in \text{Im}(\text{Fil}^r M \rightarrow \text{Fil}^r \mathcal{M}) \subseteq \hat{\text{Fil}}^r \mathcal{M}$ and hence $\text{Fil}^r \mathcal{M} = \hat{\text{Fil}}^r \mathcal{M}$. From the definition, we also can show $\phi_r = \hat{\phi}_r$. This implies the lemma. \square

Proposition 3.13. *The functor $\mathcal{M}_{\Sigma_{\infty}} : \text{Mod}_{/\Sigma_{\infty}}^{r, \phi} \rightarrow \text{Mod}_{/S_{\infty}}^{r, \phi}$ is an equivalence of categories.*

Proof. Since the functor $\mathcal{M}_{\mathfrak{S}_{\infty}}$ is an equivalence of categories for $p \geq 3$ ([9, Theorem 2.3.1]), Corollary 3.11 and Lemma 3.12 imply the proposition in this case. For $r = 0$, put $\kappa_{\Sigma} = \kappa \cap \Sigma$, where $\kappa = \text{Ker}(S \rightarrow W)$. Then, by using a natural isomorphism $\Sigma \simeq W[[u, u^{ep}/p]]$, we can show that the functor $M \mapsto M/\kappa_{\Sigma} M$ defines an equivalence of categories $\text{Mod}_{/\Sigma_{\infty}}^{0, \phi} \rightarrow \text{Mod}_{/W_{\infty}}^{\phi}$, as

in the case of the category $\text{Mod}_{/\Sigma_\infty}^{0,\phi}$. Since the diagram

$$\begin{array}{ccc} \text{Mod}_{/\Sigma_\infty}^{0,\phi} & \xrightarrow{\mathcal{M}_{\Sigma_\infty}} & \text{Mod}_{/S_\infty}^{0,\phi} \\ & \searrow & \downarrow \\ & & \text{Mod}_{/W_\infty}^\phi \end{array}$$

is commutative and the downward arrows are equivalences of categories, the proposition follows also for $p = 2$. \square

Remark 3.14. We can also define a fully faithful functor $\mathcal{M}_\Sigma : \text{Mod}_{/\Sigma}^{r,\phi} \rightarrow \text{Mod}_{/S}^{r,\phi}$ in a similar way to $\mathcal{M}_{\Sigma_\infty}$ and prove that this is an equivalence of categories. Indeed, the claim for $p \geq 3$ follows from [9, Theorem 2.2.1]. Let \mathcal{M} be in $\text{Mod}_{/S}^{0,\phi}$ and e_1, \dots, e_d be a basis of \mathcal{M} over S . Let $C \in GL_d(S)$ be the matrix such that

$$\phi(e_1, \dots, e_d) = (e_1, \dots, e_d)C.$$

Then the elements $\phi(e_1), \dots, \phi(e_d)$ also form a basis of \mathcal{M} and

$$\phi(\phi(e_1), \dots, \phi(e_d)) = (\phi(e_1), \dots, \phi(e_d))\phi(C).$$

Since $\phi(S) \subseteq \Sigma$, the Σ -module M defined by $M = \Sigma\phi(e_1) \oplus \dots \oplus \Sigma\phi(e_d)$ is stable under ϕ . Hence we see that $M \in \text{Mod}_{/\Sigma}^{0,\phi}$ and $\mathcal{M} = \mathcal{M}_\Sigma(M)$.

Proposition 3.15. *Let M be an object of $\text{Mod}_{/\Sigma_\infty}^{r,\phi}$ and set $\mathcal{M} = \mathcal{M}_{\Sigma_\infty}(M)$. Then there exists a natural isomorphism of G_{K_∞} -modules*

$$\text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, A_{\text{crys}, \infty}) \rightarrow \text{Hom}_{S, \text{Fil}^r, \phi_r}(\mathcal{M}, A_{\text{crys}, \infty}).$$

Moreover, this induces for any n an isomorphism of G_{K_n} -modules

$$\text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}})) \rightarrow \text{Hom}_{S, \text{Fil}^r, \phi_r}(\mathcal{M}, W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}})).$$

Proof. By definition, $\mathcal{M} = S \otimes_\Sigma M$ and we have a natural isomorphism

$$\text{Hom}_\Sigma(M, A_{\text{crys}, \infty}) \rightarrow \text{Hom}_S(\mathcal{M}, A_{\text{crys}, \infty}).$$

From the definition, we can check that this isomorphism induces the map in the proposition, which is injective. To prove the bijectivity, by devissage we may assume that $pM = 0$. Then both sides of this injection have the same cardinality by Lemma 3.4 and the first assertion follows. Since the sequence

$$0 \longrightarrow W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}}) \longrightarrow A_{\text{crys}, \infty} \xrightarrow{p^n} A_{\text{crys}, \infty} \longrightarrow 0$$

of the category ${}'\text{Mod}_{/\Sigma}^{r,\phi}$ is exact, the first assertion implies the second one. \square

4. A METHOD OF ABRASHKIN

In this section, we study the G_{K_n} -module $\mathrm{Hom}_{\Sigma, \mathrm{Fil}^r, \phi_r}(M, W_n^{\mathrm{PD}}(\tilde{\mathcal{O}}_{\bar{K}}))$ following Abrashkin ([3]).

Let p and $0 \leq r < p-1$ be as before. We fix a system of p -power roots of unity $\{\zeta_{p^n}\}_{n \in \mathbb{Z}_{\geq 0}}$ in \bar{K} such that $\zeta_p \neq 1$ and $\zeta_{p^n} = \zeta_{p^{n+1}}^p$ for any n , and set an element $\underline{\varepsilon}$ of R to be $(\zeta_{p^n})_{n \in \mathbb{Z}_{\geq 0}}$. Then the elements $[\underline{\varepsilon}] - 1$ and $[\underline{\varepsilon}^{1/p}] - 1$ are topologically nilpotent in $W(R)$. The element of $W(R)$

$$t = ([\underline{\varepsilon}] - 1)/([\underline{\varepsilon}^{1/p}] - 1) = 1 + [\underline{\varepsilon}^{1/p}] + [\underline{\varepsilon}^{1/p}]^2 + \cdots + [\underline{\varepsilon}^{1/p}]^{p-1}$$

is a generator of the principal ideal $\mathrm{Ker}(\theta)$. We define an element $a \in W(R)$ to be

$$a = \begin{cases} \sum_{k=1}^{p-2} p^{-1}((-1)^{p-1-k} {}_{p-1}C_k - 1)[\underline{\varepsilon}^{1/p}]^k & (p \geq 3) \\ -1 & (p = 2), \end{cases}$$

where ${}_{p-1}C_k = (p-1)!/(k!(p-1-k)!)$ is the binomial coefficient. Note that the coefficient of $[\underline{\varepsilon}^{1/p}]^k$ in each term is an integer. The element a is invertible in the ring $W(R)$, since $\theta(a) = (\zeta_p - 1)^{p-1}/p \in \mathcal{O}_{\mathbb{C}}^\times$ and the ideal $\mathrm{Ker}(\theta)$ is topologically nilpotent in $W(R)$.

The element $Z = ([\underline{\varepsilon}] - 1)^{p-1}/p$ of A_{crys} is topologically nilpotent and we have $\phi(t) = p(Z - \phi(a))$. Consider the formal power series ring $W(R)[[u']]$ with the (t, u') -adic topology and the continuous ring homomorphism $W(R)[[u']] \rightarrow A_{\mathrm{crys}}$ which sends u' to Z . Let \hat{A} denote the image of this homomorphism. Then we see that the ring \hat{A} is (t, Z) -adically complete. Since we have $Z = at^{p-1} + t^p/p$, the element t^p/p of A_{crys} is contained in the subring \hat{A} and topologically nilpotent in this subring. Hence we can consider the ring \hat{A} as a Σ -algebra by $u \mapsto [\pi]$. Put $\mathrm{Fil}^i \hat{A} = (t^i, Z)$ for $0 \leq i \leq p-1$. The Frobenius endomorphism ϕ of A_{crys} preserves \hat{A} and satisfies $\phi(\mathrm{Fil}^i \hat{A}) \subseteq p^i \hat{A}$ for $0 \leq i \leq p-1$. Set $\phi_r = p^{-r} \phi|_{\mathrm{Fil}^r \hat{A}}$. Then we can consider the ring \hat{A} also as an object of the category ${}'\mathrm{Mod}_{\Sigma}^{r, \phi}$. Put $\hat{A}_n = \hat{A}/p^n \hat{A}$ and $\hat{A}_\infty = \hat{A} \otimes_W K_0/W$. We include here a proof of the following lemma stated in [3, Subsection 3.2].

Lemma 4.1. *The natural inclusion $W(R) \rightarrow \hat{A}$ induces isomorphisms of $W(R)$ -algebras $W(R)/(([\underline{\varepsilon}] - 1)^{p-1}) \rightarrow \hat{A}/(Z)$ and $W_n(R)/(([\underline{\varepsilon}] - 1)^{p-1}) \rightarrow \hat{A}_n/(Z)$.*

Proof. For a subring B of A_{crys} , put

$$I^{[s]}B = \{x \in B \mid \phi^i(x) \in \mathrm{Fil}^s A_{\mathrm{crys}} \text{ for any } i\}$$

as in [12, Subsection 5.3]. Then we have $I^{[s]}W(R) = ([\underline{\varepsilon}] - 1)^s W(R)$ and the natural ring homomorphism

$$W(R)/I^{[s]}W(R) \rightarrow A_{\mathrm{crys}}/I^{[s]}A_{\mathrm{crys}}$$

is an injection ([12, Proposition 5.1.3, Proposition 5.3.5]). Since the element Z is contained in the ideal $I^{[p-1]}A_{\text{crys}}$, this injection factors as

$$W(R)/I^{[p-1]}W(R) \rightarrow \hat{A}/(Z) \rightarrow A_{\text{crys}}/I^{[p-1]}A_{\text{crys}}.$$

Hence the former arrow is an isomorphism and the lemma follows. \square

Therefore $\hat{A}/\text{Fil}^r \hat{A}$ is p -torsion free and $p^n \text{Fil}^r \hat{A} = \text{Fil}^r \hat{A} \cap p^n \hat{A}$. Thus we can also consider \hat{A}_n and \hat{A}_∞ as objects of the category $\text{Mod}_{\Sigma}^{r,\phi}$. The absolute Galois group G_{K_∞} acts naturally on these Σ -modules.

Lemma 4.2. *We have a natural decomposition as an R -module*

$$\hat{A}_1 = R/(t^p) \oplus (Z).$$

Proof. Consider the natural inclusion $W(R) \rightarrow \hat{A}$. We claim that this induces an injection $R/(t^p) \rightarrow \hat{A}_1$. Let x be in the ring R . If the element $[x] \in W(R)$ is contained in $p\hat{A}$, then its image in $A_{\text{crys}}/pA_{\text{crys}}$ is zero. We have an isomorphism of R -algebras

$$R[Y_1, Y_2, \dots]/(t^p, Y_1^p, Y_2^p, \dots) \rightarrow A_{\text{crys}}/pA_{\text{crys}}$$

which sends Y_i to the image of $t^{p^i}/p^i!$. Thus the element x is contained in the ideal (t^p) . Conversely, if $v_R(x) \geq p$, then we have

$$[x] = w([\underline{\varepsilon}] - 1)^{p-1} + pw'$$

for some $w, w' \in W(R)$ and this implies $[x] \in p\hat{A}$. Now we have the commutative diagram of R -algebras

$$\begin{array}{ccc} R/(t^p) & \longrightarrow & \hat{A}_1 \\ & \searrow f & \downarrow \\ & & \hat{A}_1/(Z) \end{array}$$

and the map $f : R/(t^p) \rightarrow \hat{A}_1/(Z)$ is an isomorphism by Lemma 4.1. Hence the lemma follows. \square

Since $r < p - 1$, from this lemma we can show the following lemma as in the proof of [6, Lemme 2.3.1.3].

Lemma 4.3. *The functor*

$$M \mapsto \text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, \hat{A}_\infty)$$

from $\text{Mod}_{\Sigma_\infty}^{r,\phi}$ to the category of G_{K_∞} -modules is exact.

Corollary 4.4. *For any $M \in \text{Mod}_{\Sigma_\infty}^{r,\phi}$, the natural map*

$$\text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, \hat{A}_\infty) \rightarrow \text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, A_{\text{crys}, \infty})$$

is an isomorphism of G_{K_∞} -modules. Moreover, for any n , we have an isomorphism of G_{K_∞} -modules

$$\mathrm{Hom}_{\Sigma, \mathrm{Fil}^r, \phi_r}(M, \hat{A}_n) \rightarrow \mathrm{Hom}_{\Sigma, \mathrm{Fil}^r, \phi_r}(M, A_{\mathrm{crys}}/p^n A_{\mathrm{crys}}).$$

Proof. Let us prove the first assertion. By Lemma 3.3 and Lemma 4.3, we may assume $pM = 0$. Consider the commutative diagram of rings

$$\begin{array}{ccc} \hat{A}_1 & \longrightarrow & A_{\mathrm{crys}}/pA_{\mathrm{crys}} \\ & \searrow & \downarrow \\ & & R/(t^{p-1}) \end{array}$$

whose downward arrows are defined by modulo Fil^{p-1} of the rings \hat{A}_1 and $A_{\mathrm{crys}}/pA_{\mathrm{crys}}$, respectively. Since $r < p - 1$, we have $\phi_r(\mathrm{Fil}^{p-1}\hat{A}_1) = 0$ and similarly for the ring $A_{\mathrm{crys}}/pA_{\mathrm{crys}}$. Thus these two surjections induce on the ring $R/(t^{p-1})$ the same structure of a filtered ϕ_r -module over Σ . By Corollary 3.7, we have a commutative diagram

$$\begin{array}{ccc} \mathrm{Hom}_{\Sigma, \mathrm{Fil}^r, \phi_r}(M, \hat{A}_1) & \longrightarrow & \mathrm{Hom}_{\Sigma, \mathrm{Fil}^r, \phi_r}(M, A_{\mathrm{crys}}/pA_{\mathrm{crys}}) \\ & \searrow & \downarrow \\ & & \mathrm{Hom}_{\Sigma, \mathrm{Fil}^r, \phi_r}(M, R/(t^{p-1})) \end{array}$$

whose downward arrows are isomorphisms. This concludes the proof of the first assertion. Since we have an exact sequence

$$0 \longrightarrow \hat{A}_n \longrightarrow \hat{A}_\infty \xrightarrow{p^n} \hat{A}_\infty \longrightarrow 0$$

in the category $'\mathrm{Mod}_{/\Sigma}^{r, \phi}$, the second assertion follows. \square

Since the ideal (Z) of \hat{A}_n satisfies the condition of Corollary 3.7, the Σ -algebra $\hat{A}_n/(Z)$ is naturally considered as an object of $'\mathrm{Mod}_{/\Sigma}^{r, \phi}$. We also give the ring $W_n(R)/(([\underline{\varepsilon}] - 1)^{p-1})$ the structures of a Σ -algebra and a filtered ϕ_r -module over Σ induced from those of $\hat{A}_n/(Z)$ by the isomorphism in Lemma 4.1. The map

$$\Sigma \rightarrow W_n(R)/(([\underline{\varepsilon}] - 1)^{p-1})$$

sends the element $u \in \Sigma$ to the image of $[\underline{\pi}]$ in the ring on the right-hand side. Put $v = t/E([\underline{\pi}]) \in W(R)^\times$. As for the element $Y \in \Sigma$, the equality

$$Y = -av^{-1}E([\underline{\pi}])^{p-1} + v^{-p}Z$$

holds in \hat{A} . Hence the above homomorphism sends the element Y to the image of $-av^{-1}E([\underline{\pi}])^{p-1}$.

Consider the surjective ring homomorphism

$$\begin{aligned} R &\rightarrow \tilde{\mathcal{O}}_{\tilde{K}} \\ x = (x_0, x_1, \dots) &\mapsto x_n \end{aligned}$$

and the induced surjection $\beta_n : W_n(R) \rightarrow W_n(\tilde{\mathcal{O}}_{\bar{K}})$. Let

$$J = \{(x_0, \dots, x_{n-1}) \in W_n(R) \mid v_R(x_i) \geq p^n \text{ for any } i\}$$

be the kernel of the latter surjection.

Lemma 4.5. *The ideal J is contained in the ideal $(([\underline{\varepsilon}] - 1)^{p-1})$ of the ring $W_n(R)$.*

Proof. Write the element $([\underline{\varepsilon}] - 1)^{p-1}$ also as $x = (x_0, \dots, x_{n-1}) \in W_n(R)$ with $v_R(x_0) = p$. Take an element $z = (z_0, \dots, z_{n-1})$ of the ideal J . We construct $y \in W_n(R)$ such that $xy = z$. By induction, it is enough to show that if $z_0 = \dots = z_{i-1} = 0$ for some $0 \leq i \leq n-1$ and $(x_0, \dots, x_i)(0, \dots, 0, y_i) = (0, \dots, 0, z_i)$ in $W_{i+1}(R)$, then $x(0, \dots, 0, y_i, 0, \dots, 0) \in J$. Let us write this element as $(0, \dots, 0, w_i, \dots, w_{n-1})$ with $w_i = z_i$. We have $v_R(y_i) \geq p^n - p^{i+1}$. In the ring of Witt vectors $W_n(\mathbb{F}_p[X_0, \dots, X_{n-1}, Y_0, \dots, Y_{n-1}])$, the k -th entry of the vector

$$(X_0, \dots, X_{n-1})(0, \dots, 0, Y_i, 0, \dots, 0)$$

is $X_{k-i}^{p^i} Y_i^{p^{k-i}}$ for any $k \geq i$. Thus we have $v_R(w_k) \geq p^n$. \square

Note that the elements $[\zeta_{p^n}] - 1$ and $[\zeta_{p^{n+1}}] - 1$ are nilpotent in $W_n(\tilde{\mathcal{O}}_{\bar{K}})$. By the above lemma, we have an isomorphism of rings

$$W_n(R)/(([\underline{\varepsilon}] - 1)^{p-1}) \rightarrow W_n(\tilde{\mathcal{O}}_{\bar{K}})/(([\zeta_{p^n}] - 1)^{p-1}).$$

We let $\bar{A}_{n,p-1}$ denote the ring on the right-hand side and give the ring $\bar{A}_{n,p-1}$ the structure of a filtered ϕ_r -module over Σ induced by this isomorphism.

For an algebraic extension F of K , we put

$$\mathfrak{b}_F = \{x \in \mathcal{O}_F \mid v_K(x) > er/(p-1)\}.$$

Note that the ring $\mathcal{O}_F/\mathfrak{b}_F$ is killed by p . We consider the ring of Witt vectors $W_n(\mathcal{O}_F/\mathfrak{b}_F)$ as a $W_n(\mathcal{O}_F)$ -algebra by the natural ring surjection $W_n(\mathcal{O}_F) \rightarrow W_n(\mathcal{O}_F/\mathfrak{b}_F)$ and as a W_n -algebra by twisting the natural action by σ^{-n} , as before. For a ring B and its ideal I , we define an ideal $W_n(I)$ of the ring $W_n(B)$ to be

$$W_n(I) = \{(x_0, \dots, x_{n-1}) \in W_n(B) \mid x_i \in I \text{ for any } i\}.$$

Put $F_n = K_n(\zeta_{p^{n+1}})$. For an algebraic extension F of F_n in \bar{K} , the elements $[\zeta_{p^n}] - 1$ and $[\zeta_{p^{n+1}}] - 1$ of $W_n(m_F)$ are topologically nilpotent non-zero divisors in $W_n(\mathcal{O}_F)$. Let the ring

$$W_n(\mathcal{O}_F/\mathfrak{b}_F)/([\zeta_{p^n}] - 1)^r W_n(m_F/\mathfrak{b}_F)$$

be denoted by $\bar{A}_{n,F,r+}$. We also put $\bar{A}_{n,r+} = \bar{A}_{n,\bar{K},r+}$.

Lemma 4.6. *The ideal $([\zeta_{p^n}] - 1)^r W_n(m_F)$ of $W_n(\mathcal{O}_F)$ contains the ideal $W_n(\mathfrak{b}_F)$ for any $r \in \{0, \dots, p-2\}$. We also have $(([\zeta_{p^n}] - 1)^{p-1}) \supseteq W_n(p\mathcal{O}_F)$.*

Proof. The proof is similar to the proof of Lemma 4.5. Let us show the first assertion. Since this is trivial for $r = 0$, we may assume $r \geq 1$. Put $x = (x_0, \dots, x_{n-1}) = ([\zeta_{p^n}] - 1)^r \in W_n(\mathcal{O}_F)$. Then we have $v_p(x_0) = r/(p^{n-1}(p-1))$. By induction, it is enough to show that for $0 \leq i \leq n-1$, if $(x_0, \dots, x_i)(0, \dots, 0, y_i) \in W_{i+1}(\mathfrak{b}_F)$, then $y_i \in m_F$ and $x(0, \dots, 0, y_i, 0, \dots, 0) \in W_n(\mathfrak{b}_F)$. By assumption, we have

$$v_p(y_i) > \frac{r}{p-1} \left(1 - \frac{1}{p^{n-i-1}}\right) \geq 0.$$

Put $(0, \dots, 0, w_i, \dots, w_{n-1}) = x(0, \dots, 0, y_i, 0, \dots, 0)$. We show $w_l \in \mathfrak{b}_F$ for any l by induction. Indeed, let us suppose that $w_l \in \mathfrak{b}_F$ for any $i \leq l \leq k-1$ with some $i+1 \leq k \leq n-1$. We have the equality

$$p^i y_i^{p^{k-i}} (x_0^{p^k} + p x_1^{p^{k-1}} + \dots + p^k x_k) = (p^i w_i^{p^{k-i}} + p^{i+1} w_{i+1}^{p^{k-i-1}} + \dots + p^k w_k).$$

Since $r \geq 1$, we have $(p^{k-l} - 1)r/(p-1) \geq k-l$ for $0 \leq l \leq k-1$. This implies $v_p(p^l w_l^{p^{k-l}}) > k + r/(p-1)$ for $0 \leq l \leq k-1$. The valuation of the left-hand side of the above equality also satisfies this inequality. Thus we have $v_p(w_k) > r/(p-1)$ and the assertion follows. We can show the second assertion similarly. \square

By this lemma, the natural surjections of rings

$$\begin{aligned} W_n(\mathcal{O}_F)/([\zeta_{p^n}] - 1)^r W_n(m_F) \\ \rightarrow W_n(\mathcal{O}_F/p\mathcal{O}_F)/([\zeta_{p^n}] - 1)^r W_n(m_F/p\mathcal{O}_F) \rightarrow \bar{A}_{n,F,r+} \end{aligned}$$

are isomorphisms. Then we see that the natural injection $F \rightarrow \bar{K}$ induces an injection of rings $\bar{A}_{n,F,r+} \rightarrow \bar{A}_{n,r+}$.

Write Z_n for the image of the element Z of A_{crys} in $W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}})$. Then we have a commutative diagram of Σ -algebras

$$\begin{array}{ccc} \hat{A}_n & \xrightarrow{\quad} & A_{\text{crys}}/p^n A_{\text{crys}} \\ \downarrow & & \downarrow \wr \\ & & W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}}) \\ W_n(R)/(([\varepsilon] - 1)^{p-1}) & \xrightarrow{\sim} & \hat{A}_n/(Z) \\ \downarrow \wr & \searrow & \downarrow \\ \bar{A}_{n,p-1} & \xrightarrow{\quad} & W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}})/(Z_n), \\ \downarrow & & \\ \bar{A}_{n,r+} & & \end{array}$$

where all the vertical arrows are surjections satisfying the condition of Corollary 3.7. Hence this is also a commutative diagram in $'\text{Mod}_{/\Sigma}^{r,\phi}$. Note

that these rings and homomorphisms are independent of the choice of a system $\{\zeta_{p^n}\}_{n \in \mathbb{Z}_{\geq 0}}$. We also note that $\text{Fil}^r \bar{A}_{n,r+} = E([\pi_n])^r \bar{A}_{n,r+}$ and $\phi_r(E([\pi_n])^r y) = c^r \phi(y)$ for any $y \in \bar{A}_{n,r+}$, where ϕ denotes the Frobenius endomorphism of $\bar{A}_{n,r+}$ induced from that of the ring $W_n(\mathcal{O}_{\bar{K}}/\mathfrak{b}_{\bar{K}})$. Moreover, let M be an object of $\text{Mod}_{\Sigma/\Sigma_\infty}^{r,\phi}$. Then, by Corollary 3.7 and Corollary 4.4, we have a natural isomorphism of abelian groups

$$\text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}})) \rightarrow \text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, \bar{A}_{n,r+}).$$

Next we investigate the module on the right-hand side of this isomorphism, and prove this is in fact an isomorphism of G_{F_n} -modules. Consider the element $E([\pi_n]) \in W_n(\mathcal{O}_{F_n}/p\mathcal{O}_{F_n})$ and let us fix its lift $\hat{\gamma} \in W_n(\mathcal{O}_{F_n})$ by the natural surjection $W_n(\mathcal{O}_{F_n}) \rightarrow W_n(\mathcal{O}_{F_n}/p\mathcal{O}_{F_n})$. Let $a \in W(R)^\times$ and $v = t/E([\pi]) \in W(R)^\times$ as before. We let a_n, t_n and v_n denote the images of a, t and v by the surjection $W(R) \rightarrow W_n(\tilde{\mathcal{O}}_{\bar{K}})$ induced by β_n , respectively. The elements a_n and t_n of the ring $W_n(\tilde{\mathcal{O}}_{\bar{K}})$ are contained in the subring $W_n(\mathcal{O}_{F_n}/p\mathcal{O}_{F_n})$. We abusively let them also denote their images by the natural surjections $W_n(\tilde{\mathcal{O}}_{\bar{K}}) \rightarrow W_n(\mathcal{O}_{\bar{K}}/\mathfrak{b}_{\bar{K}}) \rightarrow \bar{A}_{n,r+}$.

Lemma 4.7. *The element*

$$\hat{t}_n = 1 + [\zeta_{p^{n+1}}] + [\zeta_{p^{n+1}}]^2 + \cdots + [\zeta_{p^{n+1}}]^{p-1} = \frac{[\zeta_{p^n}] - 1}{[\zeta_{p^{n+1}}] - 1}$$

is divisible by $\hat{\gamma}$ in the ring $W_n(\mathcal{O}_{F_n})$. In particular, $\hat{\gamma}$ is a non-zero divisor of the ring $W_n(\mathcal{O}_{\bar{K}})$.

Proof. It is enough to show the divisibility in the ring $W_n(\mathcal{O}_{\bar{K}})$. Note that the element t_n is also the image of \hat{t}_n by the natural map $W_n(\mathcal{O}_{F_n}) \rightarrow W_n(\mathcal{O}_{F_n}/p\mathcal{O}_{F_n})$. Let \hat{v}_n be a lift of v_n by the natural surjection $W_n(\mathcal{O}_{\bar{K}}) \rightarrow W_n(\tilde{\mathcal{O}}_{\bar{K}})$. Then we have $\hat{t}_n - \hat{\gamma}\hat{v}_n \in W_n(p\mathcal{O}_{\bar{K}})$. By Lemma 4.6, there exists $\hat{y} \in W_n(m_{\bar{K}})$ such that $\hat{t}_n - \hat{\gamma}\hat{v}_n = \hat{t}_n\hat{y}$. Hence we have $\hat{t}_n(1 - \hat{y}) = \hat{\gamma}\hat{v}_n$. Since \hat{y} is topologically nilpotent in the ring $W_n(\mathcal{O}_{\bar{K}})$, the element $1 - \hat{y}$ is invertible and the lemma follows. \square

Lemma 4.8. *The image of $Y \in \Sigma$ in the ring $\bar{A}_{n,r+}$ (resp. $\bar{A}_{n,p-1}$) is contained in its subring $\bar{A}_{n,F_n,r+}$ (resp. $W_n(\mathcal{O}_{F_n}/p\mathcal{O}_{F_n})/([\zeta_{p^n}] - 1)^{p-1}$).*

Proof. We have the equality

$$E([\pi_n])v_n = t_n = 1 + [\zeta_{p^{n+1}}] + [\zeta_{p^{n+1}}]^2 + \cdots + [\zeta_{p^{n+1}}]^{p-1}$$

in the ring $W_n(\tilde{\mathcal{O}}_{\bar{K}})$. Note that any element $v'_n \in W_n(\tilde{\mathcal{O}}_{\bar{K}})$ satisfying the same equality is invertible and thus the elements $(v'_n)^{-1}E([\pi_n])$ are equal to each other. Since $Y = -a_nv_n^{-1}E([\pi_n])^{p-1}$ in the rings $\bar{A}_{n,r+}$ and $\bar{A}_{n,p-1}$, it suffices to construct an element v'_n of the ring $W_n(\mathcal{O}_{F_n}/p\mathcal{O}_{F_n})$ such that the equality $E([\pi_n])v'_n = t_n$ holds. This follows from Lemma 4.7. \square

From this lemma, we see that the natural G_{F_n} -actions on the rings $\bar{A}_{n,p-1}$ and $\bar{A}_{n,r+}$ are compatible with the filtered ϕ_r -module structures over Σ . In

the big commutative diagram above, the lowest horizontal arrow and lower right vertical arrow are G_K -linear by definition. Hence we have shown the following proposition.

Proposition 4.9. *Let M be an object of $\text{Mod}_{/\Sigma_\infty}^{r,\phi}$. Then the map*

$$\text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, W_n^{\text{PD}}(\tilde{\mathcal{O}}_{\bar{K}})) \rightarrow \text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, \bar{A}_{n,r+})$$

is an isomorphism of G_{F_n} -modules.

Let M be as in the proposition. Let e_1, \dots, e_d be a system of generators of M as in Lemma 3.5 and $C = (c_{i,j}) \in M_d(\Sigma)$ be a matrix representing ϕ_r as in Corollary 3.6. Consider the surjection $\Sigma^{\oplus d} \rightarrow M$ defined by $(s_1, \dots, s_d) \mapsto s_1 e_1 + \dots + s_d e_d$ and let $(s_{1,1}, \dots, s_{1,d}), \dots, (s_{q,1}, \dots, s_{q,d})$ be a system of generators of its kernel. Then the underlying G_{F_n} -set of the G_{F_n} -module

$$\text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, \bar{A}_{n,r+})$$

is identified with the set of d -tuples $(\bar{x}_1, \dots, \bar{x}_d)$ in $\bar{A}_{n,r+}$ such that the following three conditions hold:

- $s_{l,1}\bar{x}_1 + \dots + s_{l,d}\bar{x}_d = 0$ for any l ,
- $c_{1,i}\bar{x}_1 + \dots + c_{d,i}\bar{x}_d \in \text{Fil}^r \bar{A}_{n,r+}$ for any i ,
- the following equality holds:

$$\begin{cases} \phi_r(c_{1,1}\bar{x}_1 + \dots + c_{d,1}\bar{x}_d) = \bar{x}_1 \\ \vdots \\ \phi_r(c_{1,d}\bar{x}_1 + \dots + c_{d,d}\bar{x}_d) = \bar{x}_d. \end{cases}$$

We choose lifts \hat{c} , $\hat{c}_{i,j}$ and $\hat{s}_{i,j}$ in $W_n(\mathcal{O}_{F_n})$ of the images of c , $c_{i,j}$ and $s_{i,j}$ in $\bar{A}_{n,r+}$ by the natural ring homomorphism

$$W_n(\mathcal{O}_{\bar{K}}) \rightarrow W_n(\tilde{\mathcal{O}}_{\bar{K}}) \rightarrow W_n(\mathcal{O}_{\bar{K}}/\mathfrak{b}_{\bar{K}}) \rightarrow \bar{A}_{n,r+},$$

respectively. Recall that we have already chosen a lift $\hat{\gamma} \in W_n(\mathcal{O}_{F_n})$ of $E([\pi_n]) \in W_n(\mathcal{O}_{F_n}/p\mathcal{O}_{F_n})$.

Fix a polynomial $\Phi_i \in \mathbb{Z}[X_0, \dots, X_{n-1}]$ such that $\Phi_i \equiv X_i^p \pmod{p}$. This induces for any commutative ring B a map $\Phi = (\Phi_0, \dots, \Phi_{n-1}) : W_n(B) \rightarrow W_n(B)$ which is a lift of the Frobenius endomorphism on $W_n(B/pB)$. In particular, set B to be the polynomial ring $\mathbb{Z}[X_0, \dots, X_{n-1}, Y_0, \dots, Y_{n-1}]$. Put $X = (X_0, \dots, X_{n-1})$ and $Y = (Y_0, \dots, Y_{n-1})$ in the ring $W_n(B)$. Then we see that there exists elements U_0, \dots, U_{n-1} and U'_0, \dots, U'_{n-1} of the polynomial ring B such that

$$\begin{aligned} \Phi(X + Y) &= \Phi(X) + \Phi(Y) + (pU_0, \dots, pU_{n-1}), \\ \Phi(XY) &= \Phi(X)\Phi(Y) + (pU'_0, \dots, pU'_{n-1}) \end{aligned}$$

in the ring $W_n(B)$.

Proposition 4.10. *Every d -tuple $(\bar{x}_1, \dots, \bar{x}_d)$ in $\bar{A}_{n,r+}$ satisfying the above three conditions uniquely lifts to a d -tuple $(\hat{x}_1, \dots, \hat{x}_d)$ in $W_n(\mathcal{O}_{\bar{K}})$ such that*

- $\hat{s}_{l,1}\hat{x}_1 + \dots + \hat{s}_{l,d}\hat{x}_d \in ([\zeta_{p^n}] - 1)^r W_n(m_{\bar{K}})$ for any l ,

- $\hat{c}_{1,i}\hat{x}_1 + \cdots + \hat{c}_{d,i}\hat{x}_d \in \hat{\gamma}^r W_n(\mathcal{O}_{\bar{K}})$ for any i ,
- the following equality holds:

$$\begin{cases} \hat{c}^r \Phi((\hat{c}_{1,1}\hat{x}_1 + \cdots + \hat{c}_{d,1}\hat{x}_d)/\hat{\gamma}^r) = \hat{x}_1 \\ \vdots \\ \hat{c}^r \Phi((\hat{c}_{1,d}\hat{x}_1 + \cdots + \hat{c}_{d,d}\hat{x}_d)/\hat{\gamma}^r) = \hat{x}_d. \end{cases}$$

Proof. Fix a lift \hat{x}_i of \bar{x}_i in $W_n(\mathcal{O}_{\bar{K}})$. Recall that the kernel of the surjection $W_n(\mathcal{O}_{\bar{K}}) \rightarrow \bar{A}_{n,r+}$ is equal to the ideal $([\zeta_{p^n}] - 1)^r W_n(m_{\bar{K}})$. The first condition in the proposition holds automatically for $(\hat{x}_1, \dots, \hat{x}_d)$. By Lemma 4.7, the element $\hat{c}_{1,i}\hat{x}_1 + \cdots + \hat{c}_{d,i}\hat{x}_d$ is contained in $\hat{\gamma}^r W_n(\mathcal{O}_{\bar{K}})$ for any i . Since the map $\phi_r : \text{Fil}^r \bar{A}_{n,r+} \rightarrow \bar{A}_{n,r+}$ satisfies $\phi_r(E([\pi_n])^r \bar{x}) = c^r \phi(\bar{x})$ for any $\bar{x} \in \bar{A}_{n,r+}$, we have

$$\begin{cases} \hat{c}^r \Phi((\hat{c}_{1,1}\hat{x}_1 + \cdots + \hat{c}_{d,1}\hat{x}_d)/\hat{\gamma}^r) = \hat{x}_1 + ([\zeta_{p^n}] - 1)^r \hat{\delta}_1 \\ \vdots \\ \hat{c}^r \Phi((\hat{c}_{1,d}\hat{x}_1 + \cdots + \hat{c}_{d,d}\hat{x}_d)/\hat{\gamma}^r) = \hat{x}_d + ([\zeta_{p^n}] - 1)^r \hat{\delta}_d \end{cases}$$

for some $\hat{\delta}_1, \dots, \hat{\delta}_d \in W_n(m_{\bar{K}})$. It suffices to show that there exists a unique d -tuple $(\hat{y}_1, \dots, \hat{y}_d)$ in $W_n(m_{\bar{K}})$ such that

$$\begin{aligned} \hat{c}^r \Phi((\hat{c}_{1,i}(\hat{x}_1 + ([\zeta_{p^n}] - 1)^r \hat{y}_1) + \cdots + \hat{c}_{d,i}(\hat{x}_d + ([\zeta_{p^n}] - 1)^r \hat{y}_d))/\hat{\gamma}^r) \\ = \hat{x}_i + ([\zeta_{p^n}] - 1)^r \hat{y}_i \end{aligned}$$

for any i . For this, we need the following lemma.

Lemma 4.11. *Let N be a complete discrete valuation field and m_N be the maximal ideal of N . Let $\epsilon_1, \dots, \epsilon_d$ be in m_N . Let P_1, \dots, P_d and P'_1, \dots, P'_d be elements of $\mathcal{O}_N[[Y_1, \dots, Y_d]]$ such that $P_i \in (Y_1, \dots, Y_d)^2$. Then the equation*

$$\begin{cases} Y_1 - P_1(Y_1, \dots, Y_d) - \epsilon_1 P'_1(Y_1, \dots, Y_d) = 0 \\ \vdots \\ Y_d - P_d(Y_1, \dots, Y_d) - \epsilon_d P'_d(Y_1, \dots, Y_d) = 0 \end{cases}$$

has a unique solution in m_N .

Proof. By assumption, we see that for any integer $l \geq 1$, a d -tuple (y_1, \dots, y_d) in m_N/m_N^l satisfying the above equation lifts uniquely to a d -tuple in m_N/m_N^{l+1} satisfying the same equation. Thus the lemma follows. \square

Let us write as $\hat{y}_i = (\hat{y}_{i,0}, \dots, \hat{y}_{i,n-1})$. Since the image of $\Phi(([\zeta_{p^{n+1}}] - 1)^r)$ in $\bar{A}_{n,r+}$ is equal to $([\zeta_{p^n}] - 1)^r$, we can find $\hat{b} \in W_n(\mathcal{O}_{\bar{K}})$ such that

$$\Phi(([\zeta_{p^n}] - 1)^r/\hat{\gamma}^r) = ([\zeta_{p^n}] - 1)^r \hat{b}.$$

Then there exists polynomials $U_{i,m}$ over $\mathcal{O}_{\bar{K}}$ of the indeterminates $\underline{Y} = (Y_{i,m})_{1 \leq i \leq d, 0 \leq m \leq n-1}$ such that the equation we have to solve is

$$\begin{aligned} \hat{x}_i + ([\zeta_{p^n}] - 1)^r \hat{y}_i &= \hat{x}_i + ([\zeta_{p^n}] - 1)^r \hat{\delta}_i \\ &\quad + ([\zeta_{p^n}] - 1)^r \hat{b} \hat{c}^r (\Phi(\hat{c}_{1,i})\Phi(\hat{y}_1) + \cdots + \Phi(\hat{c}_{d,i})\Phi(\hat{y}_d)) \\ &\quad + (pU_{i,0}(\underline{\hat{y}}), \dots, pU_{i,n-1}(\underline{\hat{y}})) \end{aligned}$$

for any i , where we put $\underline{\hat{y}} = (\hat{y}_{i,m})_{1 \leq i \leq d, 0 \leq m \leq n-1}$. As in the proof of Lemma 4.6, we see that, for any elements P_0, \dots, P_{n-1} of the polynomial ring $\mathcal{O}_{\bar{K}}[\underline{Y}]$, we can uniquely find elements Q_0, \dots, Q_{n-1} of this ring such that the coefficients of these polynomials are in the maximal ideal $m_{\bar{K}}$ and the equality

$$(pP_0, \dots, pP_{n-1}) = ([\zeta_{p^n}] - 1)^r (Q_0, \dots, Q_{n-1})$$

holds in the ring of Witt vectors $W_n(\mathcal{O}_{\bar{K}}[\underline{Y}])$. Therefore, this equation is equivalent to the equation

$$\begin{aligned} \hat{y}_i &= \hat{\delta}_i + \hat{b} \hat{c}^r (\Phi(\hat{c}_{1,i})\Phi(\hat{y}_1) + \cdots + \Phi(\hat{c}_{d,i})\Phi(\hat{y}_d)) \\ &\quad + (V_{i,0}(\underline{\hat{y}}), \dots, V_{i,n-1}(\underline{\hat{y}})), \end{aligned}$$

where $V_{i,m}$ is a polynomial of \underline{Y} over $\mathcal{O}_{\bar{K}}$ whose coefficients are in the maximal ideal $m_{\bar{K}}$. From the definition of Φ , we see that $\underline{\hat{y}} = (\hat{y}_{i,m})_{i,m}$ is a solution of a system of equations

$$Y_{i,m} - P_{i,m}(\underline{Y}) - \epsilon_{i,m} P'_{i,m}(\underline{Y}) = 0$$

satisfying the condition of Lemma 4.11 for a sufficiently large finite extension N of K . Then, by this lemma, we can solve the equation uniquely in $m_{\bar{K}}$. \square

Let F be an algebraic extension of F_n in \bar{K} and consider the ring $\bar{A}_{n,F,r+}$. By Lemma 4.8, we can consider this ring as a Σ -subalgebra of $\bar{A}_{n,r+}$. Put $\text{Fil}^r \bar{A}_{n,F,r+} = E([\pi_n])^r \bar{A}_{n,F,r+}$. Then Lemma 4.7 implies that

$$\bar{A}_{n,F,r+} \cap \text{Fil}^r \bar{A}_{n,r+} = \text{Fil}^r \bar{A}_{n,F,r+}.$$

Moreover, the Frobenius endomorphism ϕ of the ring $\bar{A}_{n,r+}$ preserves the subalgebra $\bar{A}_{n,F,r+}$ and thus $\phi_r : \text{Fil}^r \bar{A}_{n,r+} \rightarrow \bar{A}_{n,r+}$ induces a ϕ -semilinear map $\phi_r : \text{Fil}^r \bar{A}_{n,F,r+} \rightarrow \bar{A}_{n,F,r+}$. Hence $\bar{A}_{n,F,r+}$ is a subobject of $\bar{A}_{n,r+}$ in the category ${}'\text{Mod}_{/\Sigma}^{r,\phi}$. For $M \in \text{Mod}_{/\Sigma_\infty}^{r,\phi}$, let us set

$$T_{\text{crys}, \pi_n, F}^*(M) = \text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, \bar{A}_{n,F,r+}).$$

We see that

$$\bar{A}_{n,r+} = \bar{A}_{n,\bar{K},r+} = \bigcup_{F/F_n} \bar{A}_{n,F,r+}$$

in ${}'\text{Mod}_{/\Sigma}^{r,\phi}$ and thus we have a natural identification of abelian groups

$$T_{\text{crys}, \pi_n, \bar{K}}^*(M) = \bigcup_{F/F_n} T_{\text{crys}, \pi_n, F}^*(M).$$

The absolute Galois group G_{F_n} acts on the abelian group on the left-hand side.

Lemma 4.12. *Let F be an algebraic extension of F_n in \bar{K} . Then the G_F -fixed part $T_{\text{crys}, \pi_n, \bar{K}}^*(M)^{G_F}$ is equal to $T_{\text{crys}, \pi_n, F}^*(M)$.*

Proof. From Proposition 4.10, we see that the elements of $T_{\text{crys}, \pi_n, \bar{K}}^*(M)$ correspond bijectively to the d -tuples in $W_n(\mathcal{O}_{\bar{K}})$ satisfying the three conditions in this proposition. The uniqueness assertion of the proposition shows that $g \in G_F$ fixes such a d -tuple in $W_n(\mathcal{O}_{\bar{K}})$ if and only if g fixes its image in $\bar{A}_{n, r+}$. Hence an element of $T_{\text{crys}, \pi_n, \bar{K}}^*(M)$ is fixed by G_F if and only if it is contained in the image of $W_n(\mathcal{O}_F)$. Thus the lemma follows. \square

Corollary 4.13. *Let L_n be the finite Galois extension of F_n corresponding to the kernel of the map*

$$G_{F_n} \rightarrow \text{Aut}(T_{\text{crys}, \pi_n, \bar{K}}^*(M)).$$

Then an algebraic extension F of F_n in \bar{K} contains L_n if and only if

$$\#T_{\text{crys}, \pi_n, F}^*(M) = \#T_{\text{crys}, \pi_n, \bar{K}}^*(M).$$

Proof. An algebraic extension F of F_n contains L_n if and only if the action of G_F on $T_{\text{crys}, \pi_n, \bar{K}}^*(M)$ is trivial. By Lemma 4.12, this is equivalent to $T_{\text{crys}, \pi_n, F}^*(M) = T_{\text{crys}, \pi_n, \bar{K}}^*(M)$. \square

5. RAMIFICATION BOUND

In this section, we prove Theorem 1.1. Let \mathcal{M} be an object of $\text{Mod}_{/S_\infty}^{r, \phi, N}$ which is killed by p^n and let L be the finite Galois extension of K corresponding to the kernel of the map

$$G_K \rightarrow \text{Aut}(T_{\text{st}, \pi}^*(\mathcal{M})).$$

Then the theorem is equivalent to the inequality $u_{L/K} \leq u(K, r, n)$, where $u_{L/K}$ denotes the greatest upper ramification break of the Galois extension L/K ([10]). For $r = 0$, the G_K -module $T_{\text{st}, \pi}^*(\mathcal{M})$ is unramified and the assertion is trivial. Thus we may assume $p \geq 3$ and $r \geq 1$.

Let L_n be the finite Galois extension of F_n corresponding to the kernel of the map

$$G_{F_n} \rightarrow \text{Aut}(T_{\text{st}, \pi}^*(\mathcal{M})).$$

Since F_n is Galois over K , the extension $L_n = LF_n$ is also a Galois extension of K . Let $M \in \text{Mod}_{/\Sigma_\infty}^{r, \phi}$ be the filtered ϕ_r -module over Σ which corresponds to \mathcal{M} by the equivalence $\mathcal{M}_{\Sigma_\infty}$ of Proposition 3.13. Then Proposition 3.15 and Proposition 4.9 show that L_n is also the finite extension of F_n cut out by the G_{F_n} -module $T_{\text{crys}, \pi_n, \bar{K}}^*(M)$. It is enough to prove the inequality $u_{L_n/K} \leq u(K, r, n)$.

Before proving this, we state some general lemmas to calculate the ramification bound. Let N be a complete discrete valuation field of positive residue characteristic, v_N be its valuation normalized as $v_N(N^\times) = \mathbb{Z}$ and N^{sep} be its separable closure. We extend v_N to any algebraic closure of N .

Lemma 5.1. *Let $f(T) \in \mathcal{O}_N[T]$ be a separable monic polynomial and z_1, \dots, z_d be the zeros of f in $\mathcal{O}_{N^{\text{sep}}}$. Suppose that the set $\{v_N(z_k - z_i) \mid k = 1, \dots, d, k \neq i\}$ is independent of i . Put*

$$s_f = \sum_{\substack{k=1, \dots, d \\ k \neq i}} v_N(z_k - z_i) \text{ and } \alpha_f = \sup_{\substack{k=1, \dots, d \\ k \neq i}} v_N(z_k - z_i),$$

which are independent of i by assumption. If $j > s_f + \alpha_f$, then we have the decomposition

$$\{x \in \mathcal{O}_{N^{\text{sep}}} \mid v_N(f(x)) \geq j\} = \coprod_{i=1, \dots, d} \{x \in \mathcal{O}_{N^{\text{sep}}} \mid v_N(x - z_i) \geq j - s_f\}.$$

Otherwise, the set on the left-hand side contains

$$\{x \in \mathcal{O}_{N^{\text{sep}}} \mid v_N(x - z_i) \geq \alpha_f\},$$

which contains at least two zeros of f .

Proof. A verbatim argument in the proof of [1, Lemma 6.6] shows the claim. \square

Corollary 5.2. *Let $f(T)$ be as above and put $B = \mathcal{O}_N[T]/(f(T))$. Let us write the N -algebra $N' = B \otimes_{\mathcal{O}_N} N$ as the product $N_1 \times \dots \times N_t$ of finite separable extensions N_1, \dots, N_t of N . If $j > s_f + \alpha_f$, then the j -th upper numbering ramification group ([1]), which we let be denoted by $G_N^{(j)}$, is contained in G_{N_i} for any i . Moreover, if N' is a field and B coincides with $\mathcal{O}_{N'}$, then $j > s_f + \alpha_f$ if and only if $G_N^{(j)} \subseteq G_{N'}$.*

Proof. Note that the algebra B is finite flat and of relative complete intersection over \mathcal{O}_N . By the previous lemma, the conductor $c(B)$ of the \mathcal{O}_N -algebra B ([1, Proposition 6.4]) is equal to $s_f + \alpha_f$. Thus we have the inequality

$$c(\mathcal{O}_{N_1} \times \dots \times \mathcal{O}_{N_t}) \leq c(B) = s_f + \alpha_f$$

by the definition of the conductor and a functoriality of the functor \mathcal{F}^j defined in [1]. This implies the corollary. \square

Corollary 5.3. *We have the inequality*

$$u_{K(\zeta_{p^{n+1}})/K} \leq 1 - \frac{1}{e(K(\zeta_p)/K)} + e(n + \frac{1}{p-1}),$$

where $e(K(\zeta_p)/K)$ denotes the relative ramification index of $K(\zeta_p)$ over K .

Proof. Since the Herbrand function is transitive and the finite extension $K(\zeta_p)$ is tamely ramified over K , it is enough to show the inequality

$$u_{K(\zeta_{p^{n+1}})/K(\zeta_p)} \leq e(K(\zeta_p))(n + \frac{1}{p-1}).$$

Put $N = K(\zeta_p)$ and $f(T) = T^{p^n} - \zeta_p$. These satisfy the assumptions of Corollary 5.2. We have $s_f = ne(K(\zeta_p))$ and $\alpha_f = e(K(\zeta_p))/(p-1)$ in this case. Hence the corollary follows. \square

Corollary 5.4. *Consider the finite Galois extension $F_n = K_n(\zeta_{p^{n+1}})$ of K . Then we have the equality*

$$u_{F_n/K} = 1 + e(n + \frac{1}{p-1}).$$

Proof. Applying Corollary 5.2 to the Eisenstein polynomial $f(T) = T^{p^n} - \pi$ and $N = K$ shows that $j > 1 + e(n + 1/(p-1))$ if and only if $G_K^{(j)} \subseteq G_{K_n}$. From Corollary 5.3, we see that if $j > 1 + e(n + 1/(p-1))$, then $G_K^{(j)} \subseteq G_{K(\zeta_{p^{n+1}})}$. Since $G_{F_n} = G_{K_n} \cap G_{K(\zeta_{p^{n+1}})}$, we conclude that $j > 1 + e(n + 1/(p-1))$ if and only if $G_K^{(j)} \subseteq G_{F_n}$. \square

Remark 5.5. Note that this argument also shows the equality

$$u_{K_n(\zeta_{p^n})/K} = 1 + e(n + \frac{1}{p-1}).$$

Next we assume that the residue field of N is perfect. For an algebraic extension F of N , we put

$$\mathfrak{a}_{F/N}^j = \{x \in \mathcal{O}_F \mid v_N(x) \geq j\}.$$

Let Q be a finite Galois extension of N and consider the property

$$(P_j) \begin{cases} \text{for any algebraic extension } F \text{ of } N, \text{ if there exists} \\ \text{an } \mathcal{O}_N\text{-algebra homomorphism } \mathcal{O}_Q \rightarrow \mathcal{O}_F/\mathfrak{a}_{F/N}^j, \\ \text{then there exists an } N\text{-algebra injection } Q \rightarrow F \end{cases}$$

for $j \in \mathbb{R}_{\geq 0}$, as in [10, Proposition 1.5]. Then we have the following proposition, which is due to Yoshida. Here we reproduce his proof for the convenience of the reader.

Proposition 5.6 ([19]).

$$u_{Q/N} = \inf\{j \in \mathbb{R}_{\geq 0} \mid \text{the property } (P_j) \text{ holds}\}.$$

Proof. By [10, Proposition 1.5 (i)], it is enough to show that the property (P_j) does not hold for $j = u_{Q/N} - (e')^{-1}$ with an arbitrarily large $e' > 0$. As in the proof of [10, Proposition 1.5 (ii)], we may assume that Q is totally and wildly ramified over N . Take an arbitrarily large integer $e'' > 0$ with $(e'', pe(Q/N)) = 1$. We may also assume that N contains a primitive e'' -th root of unity. Set $N' = N(\pi_N^{1/e''})$ and $Q' = QN'$. Note that we have $u_{Q'/N} = u_{Q/N}$ by assumption. From this proposition in [10], we see that for some algebraic extension F of N , there exists an \mathcal{O}_N -algebra homomorphism $\mathcal{O}_{Q'} \rightarrow \mathcal{O}_F/\mathfrak{a}_{F/N}^j$ for $j = u_{Q/N} - e(Q'/N)^{-1}$ but no N -algebra injection $Q' \rightarrow F$. Since Q/N is wildly ramified, we see that $e(Q/N)u_{Q/N} - 1 > e(Q/N)$. Hence we have $u_{Q/N} - e(Q'/N)^{-1} > 1 \geq u_{N'/N}$ and there exists an N -algebra injection $N' \rightarrow F$ also by this proposition. Thus there exists no N -algebra injection $Q \rightarrow F$ and the property (P_j) for Q/N does not hold. Since $e(Q'/N) = e''e(Q/N)$, the proposition follows. \square

We see from Proposition 5.6 that to bound the greatest upper ramification break $u_{L_n/K}$, it is enough to show the following proposition.

Proposition 5.7. *Let F be an algebraic extension of K . If $j > u(K, r, n)$ and there exists an \mathcal{O}_K -algebra homomorphism*

$$\eta : \mathcal{O}_{L_n} \rightarrow \mathcal{O}_F / \mathfrak{a}_{F/K}^j,$$

then there exists a K -algebra injection $L_n \rightarrow F$.

Proof. We may assume that F is contained in \bar{K} . By assumption, we have $j > er/(p-1)$ and we see that the ideal $\mathfrak{b}_F = \{x \in \mathcal{O}_F \mid v_K(x) > er/(p-1)\}$ contains $\mathfrak{a}_{F/K}^j$. Thus η induces an \mathcal{O}_K -algebra homomorphism

$$\mathcal{O}_{L_n} \rightarrow \mathcal{O}_F / \mathfrak{b}_F.$$

Since η also induces an \mathcal{O}_K -algebra homomorphism $\mathcal{O}_{F_n} \rightarrow \mathcal{O}_F / \mathfrak{a}_{F/K}^j$ and $r \geq 1$, from Corollary 5.4 and [10, Proposition 1.5] we get a K -linear injection $F_n \rightarrow F$. Thus we see that F contains π_n and $\zeta_{p^{n+1}}$. More precisely, we have the following two lemmas.

Lemma 5.8. *There exists $i \in \mathbb{Z}$ such that $\eta(\pi_n) \equiv \pi_n \zeta_{p^n}^i \pmod{\mathfrak{b}_F}$.*

Proof. Since the map η is \mathcal{O}_K -linear, the equality $\eta(\pi_n)^{p^n} = \pi$ holds in $\mathcal{O}_F / \mathfrak{a}_{F/K}^j$. Set \hat{x} to be a lift of $\eta(\pi_n)$ in \mathcal{O}_F . Then we have

$$v_K(\hat{x}^{p^n} - \pi) = \sum_{i=0}^{p^n-1} v_K(\hat{x} - \pi_n \zeta_{p^n}^i) \geq j.$$

Let us apply Lemma 5.1 to $f(T) = T^{p^n} - \pi \in \mathcal{O}_K[T]$. Then, with the notation of the lemma, we have

$$s_f = 1 - \frac{1}{p^n} + ne \text{ and } \alpha_f = \frac{1}{p^n} + \frac{e}{p-1}.$$

Since $j - s_f > er/(p-1)$ by assumption, we have

$$\hat{x} \equiv \pi_n \zeta_{p^n}^i \pmod{\mathfrak{b}_F}$$

for some i . □

Lemma 5.9. *There exists $g' \in G_K$ such that $\eta(\zeta_{p^{n+1}}) \equiv g'(\zeta_{p^{n+1}}) \pmod{\mathfrak{b}_F}$.*

Proof. Set N to be the maximal unramified subextension of $K(\zeta_{p^{n+1}})/K$. Since the map $\mathcal{O}_K \rightarrow \mathcal{O}_N$ is etale, there exists a K -algebra injection $g_0 : N \rightarrow F$ such that $\eta(x) \equiv g_0(x) \pmod{\mathfrak{a}_{F/K}^j}$ for any $x \in \mathcal{O}_N$. Let ϖ be a uniformizer of $K(\zeta_{p^{n+1}})$ and $f(T) \in \mathcal{O}_N[T]$ be the Eisenstein polynomial of ϖ over \mathcal{O}_N . We let $f^{g_0}(T) \in \mathcal{O}_N[T]$ denote the conjugate of f by g_0 . Then f^{g_0} satisfies the conditions of Lemma 5.1. By definition we have $s_{f^{g_0}} = s_f$ and $\alpha_{f^{g_0}} = \alpha_f$. Since the roots of $f^{g_0}(T)$ are conjugates of ϖ over K , Lemma 5.1 implies as in the previous lemma that there exists $g' \in G_K$ such

that $g'|_N = g_0$ and $\eta(\varpi) \equiv g'(\varpi) \bmod \mathfrak{a}_{F/K}^{j-s_f}$. Since $\mathcal{O}_{K(\zeta_{p^{n+1}})}$ is generated by ϖ over \mathcal{O}_N , we see that $\eta(\zeta_{p^{n+1}}) \equiv g'(\zeta_{p^{n+1}}) \bmod \mathfrak{a}_{F/K}^{j-s_f}$.

Thus it is enough to check the inequality $j - s_f > er/(p-1)$. Note that s_f is equal to the valuation $v_K(\mathfrak{D}_{K(\zeta_{p^{n+1}})/N})$ of the different of the totally ramified Galois extension $K(\zeta_{p^{n+1}})/N$. To bound this, put $G = \text{Gal}(K(\zeta_{p^{n+1}})/N(\zeta_p))$ and $e' = e(N(\zeta_p)/N)$. We have

$$v_K(\tau(\varpi) - \varpi) \leq v_K(\tau(\zeta_{p^{n+1}}) - \zeta_{p^{n+1}})$$

for any $\tau \in G$ and thus

$$v_K(\mathfrak{D}_{K(\zeta_{p^{n+1}})/N(\zeta_p)}) \leq \sum_{\tau \neq 1 \in G} v_K(\tau(\zeta_{p^{n+1}}) - \zeta_{p^{n+1}}) \leq ne.$$

We also have the equality $v_K(\mathfrak{D}_{N(\zeta_p)/N}) = 1 - 1/e'$ and hence we get

$$s_f = v_K(\mathfrak{D}_{K(\zeta_{p^{n+1}})/N}) \leq 1 - 1/e' + ne.$$

Since $e' \leq p-1$, the inequality $j - s_f > er/(p-1)$ holds. \square

Corollary 5.10. *There exists $g \in G_K$ such that $\eta(\pi_n) \equiv g(\pi_n) \bmod \mathfrak{b}_F$ and $\eta(\zeta_{p^{n+1}}) \equiv g(\zeta_{p^{n+1}}) \bmod \mathfrak{b}_F$.*

Proof. Let $i \in \mathbb{Z}$ and $g' \in G_K$ be as in Lemma 5.8 and Lemma 5.9, respectively. Since $K_n \cap K(\zeta_{p^{n+1}}) = K$ (see for example [17, Lemma 5.1.2]), we can find an element $g \in G_K$ such that $g(\pi_n) = \pi_n \zeta_{p^n}^i$ and $g(\zeta_{p^{n+1}}) = g'(\zeta_{p^{n+1}})$. \square

Lemma 5.11. *For $m \in \mathbb{Z}_{\geq 0}$, set an ideal $\mathfrak{b}_{L_n}^{(m)}$ of \mathcal{O}_{L_n} to be*

$$\mathfrak{b}_{L_n}^{(m)} = \{x \in \mathcal{O}_{L_n} \mid v_K(x) > \frac{er}{p^m(p-1)}\}$$

and similarly for F . Then the \mathcal{O}_K -algebra homomorphism η induces an \mathcal{O}_K -algebra injection

$$\eta^{(m)} : \mathcal{O}_{L_n}/\mathfrak{b}_{L_n}^{(m)} \rightarrow \mathcal{O}_F/\mathfrak{b}_F^{(m)}$$

for any m .

Proof. We may assume that L_n is totally ramified over K . We write the Eisenstein polynomial of a uniformizer π_{L_n} of L_n over \mathcal{O}_K as

$$P(T) = T^{e'} + c_1 T^{e'-1} + \cdots + c_{e'-1} T + c_{e'},$$

where $e' = e(L_n/K)$. Then $z = \eta(\pi_{L_n})$ satisfies $P(z) = 0$ in $\mathcal{O}_F/\mathfrak{a}_{F/K}^j$. Let \hat{z} be a lift of z in \mathcal{O}_F . Since $j > 1$, we have $v_K(\hat{z}) = 1/e'$. The condition $i > e(L_n)r/(p^m(p-1))$ is equivalent to the condition

$$v_K(\hat{z}^i) > \frac{e(L_n)r}{p^m(p-1)} \cdot \frac{1}{e'} = \frac{er}{p^m(p-1)}.$$

Thus the claim follows. \square

Since L_n contains F_n , we can consider the ring

$$\bar{A}_{n,L_n,r+} = W_n(\mathcal{O}_{L_n}/\mathfrak{b}_{L_n})/([\zeta_{p^n}] - 1)^r W_n(m_{L_n}/\mathfrak{b}_{L_n})$$

and similarly $\bar{A}_{n,F,r+}$ for F . We give these rings structures of Σ -algebras as follows. The ring $\bar{A}_{n,L_n,r+}$ is considered as a Σ -algebra by using the system $\{\pi_n\}_{n \in \mathbb{Z}_{\geq 0}}$ we chose of p -power roots of π , as in the previous section. On the other hand, using $g \in G_K$ in Corollary 5.10, put $\tilde{\pi}_n = g(\pi_n)$ and $\tilde{\zeta}_{p^{n+1}} = g(\zeta_{p^{n+1}})$. Then we consider the ring $\bar{A}_{n,F,r+}$ as a Σ -algebra by using a system of p -power roots of π containing $\tilde{\pi}_n$. We define Fil^r and ϕ_r of these rings in the same way as before.

Lemma 5.12. *The induced ring homomorphism*

$$\bar{\eta} : \bar{A}_{n,L_n,r+} \rightarrow \bar{A}_{n,F,r+}$$

is a morphism of the category $'\text{Mod}'_{\Sigma}^{r,\phi}$.

Proof. Firstly, we check that $\bar{\eta}$ is Σ -linear. By definition, this homomorphism commutes with the action of the element $u \in \Sigma$. To show the compatibility with the element $Y \in \Sigma$, let us consider the commutative diagram

$$\begin{array}{ccc} W_n(\mathcal{O}_{L_n}/\mathfrak{b}_{L_n}) & \xrightarrow{\eta_n} & W_n(\mathcal{O}_F/\mathfrak{b}_F) \\ \downarrow & & \downarrow \\ \bar{A}_{n,L_n,r+} & \xrightarrow{\bar{\eta}} & \bar{A}_{n,F,r+}, \end{array}$$

where the horizontal arrows are induced by η . Note that we have $\eta_n([\pi_n]) = [\tilde{\pi}_n]$ and $\eta_n([\zeta_{p^{n+1}}]) = [\tilde{\zeta}_{p^{n+1}}]$. Let $a \in W(R)^\times$ and $v = t/E([\pi]) \in W(R)^\times$ be as in the previous section. Let a_n and v_n denote the images of a and v in $W_n(\mathcal{O}_{L_n}/\mathfrak{b}_{L_n})$, respectively. Then the element v_n is a solution of the equation

$$E([\pi_n])v_n = 1 + [\zeta_{p^{n+1}}] + \cdots + [\zeta_{p^{n+1}}]^{p-1}.$$

Similarly, we define elements \tilde{a}_n and \tilde{v}_n of $W_n(\mathcal{O}_F/\mathfrak{b}_F)$ using $\tilde{\pi}_n$ and $\tilde{\zeta}_{p^{n+1}}$. By definition, the element \tilde{v}_n is a solution of the equation

$$E([\tilde{\pi}_n])\tilde{v}_n = 1 + [\tilde{\zeta}_{p^{n+1}}] + \cdots + [\tilde{\zeta}_{p^{n+1}}]^{p-1}.$$

Now what we have to show is the equality

$$\bar{\eta}(a_n v_n^{-1} E([\pi_n])^{p-1}) = \tilde{a}_n \tilde{v}_n^{-1} E([\tilde{\pi}_n])^{p-1}$$

in the ring $\bar{A}_{n,F,r+}$. Since the element a_n of $W_n(\mathcal{O}_{L_n}/\mathfrak{b}_{L_n})$ is a linear combination of the elements $1, [\zeta_{p^{n+1}}], \dots, [\zeta_{p^{n+1}}]^{p-1}$ over \mathbb{Z} , we have $\bar{\eta}(a_n) = \tilde{a}_n$ in $\bar{A}_{n,F,r+}$. The elements \tilde{v}_n and $\bar{\eta}(v_n)$ satisfy the same equation in $\bar{A}_{n,F,r+}$. Since these two elements are invertible, we get $\bar{\eta}(v_n)^{-1} E([\tilde{\pi}_n]) = \tilde{v}_n^{-1} E([\tilde{\pi}_n])$ and the equality holds. Since the diagram above is compatible with the Frobenius endomorphisms, we see from the definition that $\bar{\eta}$ also preserves Fil^r and commutes with ϕ_r of both sides. \square

Thus we get a homomorphism of abelian groups

$$T_{\text{crys}, L_n, \pi_n}^*(M) \rightarrow T_{\text{crys}, F, \tilde{\pi}_n}^*(M).$$

Then the following lemma, whose proof is omitted in [3, Subsection 3.13], implies that this homomorphism is an injection. We insert here a proof of this lemma for the convenience of the reader.

Lemma 5.13. *The ring homomorphism $\bar{\eta} : \bar{A}_{n, L_n, r+} \rightarrow \bar{A}_{n, F, r+}$ is an injection.*

Proof. Let $x = (x_0, \dots, x_{n-1})$ be an element of $W_n(\mathcal{O}_{L_n}/\mathfrak{b}_{L_n})$ such that

$$(\eta^{(0)}(x_0), \dots, \eta^{(0)}(x_{n-1})) \in ([\zeta_{p^n}] - 1)^r W_n(m_F/\mathfrak{b}_F),$$

where $\eta^{(0)}$ is as in Lemma 5.11. Suppose that $x_0 = \dots = x_{m-1} = 0$ for some $0 \leq m \leq n-1$. Let $\hat{z}_i \in \mathcal{O}_F$ be a lift of $\eta^{(0)}(x_i)$. By Lemma 4.6, we have

$$(0, \dots, 0, \hat{z}_m, \dots, \hat{z}_{n-1}) = ([\zeta_{p^n}] - 1)^r (\hat{y}_0, \dots, \hat{y}_{n-1})$$

for some $\hat{y}_0, \dots, \hat{y}_{n-1} \in m_F$. Thus we get $\hat{y}_0 = \dots = \hat{y}_{m-1} = 0$ and $v_K(\hat{z}_m) > er/(p^{n-1-m}(p-1))$. Then Lemma 5.11 implies that x_m is contained in the ideal $\mathfrak{b}_{L_n}^{(n-1-m)}/\mathfrak{b}_{L_n}$ and

$$x = ([\zeta_{p^n}] - 1)^r (0, \dots, 0, y, 0, \dots, 0) + (0, \dots, 0, x'_{m+1}, \dots, x'_{n-1})$$

for some $y \in m_{L_n}/\mathfrak{b}_{L_n}$ and $x'_{m+1}, \dots, x'_{n-1} \in \mathcal{O}_{L_n}/\mathfrak{b}_{L_n}$. Repeating this, we see that x is zero in $\bar{A}_{n, L_n, r+}$ and the lemma follows. \square

Now Corollary 4.13 shows that the abelian group $T_{\text{crys}, L_n, \pi_n}^*(M)$ has the same cardinality as $T_{\text{crys}, \bar{K}, \pi_n}^*(M)$. This implies that the abelian group $T_{\text{crys}, F, \tilde{\pi}_n}^*(M)$ has cardinality no less than $\#T_{\text{crys}, \bar{K}, \pi_n}^*(M)$. Let $g \in G_K$ be as in Corollary 5.10. Then we have the following lemma.

Lemma 5.14. *The G_{F_n} -module $T_{\text{crys}, \bar{K}, \tilde{\pi}_n}^*(M)$ is isomorphic to the conjugate of the G_{F_n} -module $T_{\text{crys}, \bar{K}, \pi_n}^*(M)$ by the element g .*

Proof. Let us consider the composite

$$\Sigma \rightarrow \bar{A}_{n, r+} \xrightarrow{g} \bar{A}_{n, r+}$$

of the ring homomorphism defined by $u \mapsto [\pi_n]$ and the map induced by g . We can check that this is the natural ring homomorphism defined by $u \mapsto [\tilde{\pi}_n]$ as in the proof of Lemma 5.12. Thus we have an isomorphism of abelian groups

$$\begin{aligned} \text{Hom}_{\Sigma}(M, \bar{A}_{n, r+}) &\rightarrow \text{Hom}_{\Sigma}(M, \bar{A}_{n, r+}) \\ f &\mapsto g \circ f, \end{aligned}$$

where we consider on the ring $\bar{A}_{n, r+}$ on the right-hand side the filtered ϕ_r -module structure over Σ defined by $\tilde{\pi}_n$. As in the proof of Lemma 5.12, we can check that this isomorphism induces an injection

$$\text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, \bar{A}_{n, r+}) \rightarrow \text{Hom}_{\Sigma, \text{Fil}^r, \phi_r}(M, \bar{A}_{n, r+}).$$

This is also an isomorphism, for the map $f \mapsto g^{-1} \circ f$ defines its inverse. \square

Thus we have $\#T_{\text{crys}, \bar{K}, \tilde{\pi}_n}^*(M) = \#T_{\text{crys}, \bar{K}, \pi_n}^*(M)$. Since L_n is Galois over K , this lemma also shows that the finite Galois extension of F_n cut out by the action on $T_{\text{crys}, \bar{K}, \tilde{\pi}_n}^*(M)$ is L_n . Hence we see from Corollary 4.13 that F contains L_n and Proposition 5.7 follows. This concludes the proof of Theorem 1.1. \square

Proof of Corollary 1.3. The second assertion follows immediately from Theorem 1.1 and [8, Théorème 1.1]. As for the first assertion, note that if $r = 0$ then V is unramified and the assertion is trivial. Thus we may assume $p \geq 3$. Since we have the natural surjection $\mathcal{L}/p^n\mathcal{L} \rightarrow \mathcal{L}/\mathcal{L}'$, we may also assume $\mathcal{L}' = p^n\mathcal{L}$. For $\hat{\mathcal{M}} \in \text{Mod}_{/S}^{r, \phi, N}$, let us consider the G_K -module

$$T_{\text{st}, \underline{\pi}}^*(\hat{\mathcal{M}}) = \text{Hom}_{S, \text{Fil}^r, \phi_r, N}(\hat{\mathcal{M}}, \hat{A}_{\text{st}}).$$

By [17, Theorem 2.3.5], there exists $\hat{\mathcal{M}} \in \text{Mod}_{/S}^{r, \phi, N}$ such that the G_K -module \mathcal{L} is isomorphic to $T_{\text{st}, \underline{\pi}}^*(\hat{\mathcal{M}})$. Then we see that the G_K -module $\mathcal{L}/p^n\mathcal{L}$ is isomorphic to $T_{\text{st}, \underline{\pi}}^*(\hat{\mathcal{M}}/p^n\hat{\mathcal{M}})$ and the assertion follows from Theorem 1.1. \square

Remark 5.15. The ramification bound in Theorem 1.1 is sharp for $r \leq 1$. Indeed, the greatest upper ramification break $1 + e(n + 1/(p - 1))$ for $r = 1$ is obtained by the p^n -torsion of the Tate curve $\bar{K}^\times/\pi^\mathbb{Z}$ (see Remark 5.5). The author does not know whether these bounds are sharp also for $r \geq 2$.

REFERENCES

- [1] A. Abbes and T. Saito: *Ramification of local fields with imperfect residue fields I*, Amer. J. Math. **124** (2002), 879–920.
- [2] V. Abrashkin: *Modular representations of the Galois group of a local field, and a generalization of the Shafarevich conjecture*, Math. USSR Izvestiya **35** (1990), no. 3, 469–518.
- [3] V. Abrashkin: *Ramification in étale cohomology*, Invent. Math. **101** (1990), no. 3, 631–640.
- [4] C. Breuil: *Construction de représentations p -adiques semi-stables*, Ann. Sci. École Norm. Sup. (4) **31** (1998), no. 3, 281–327.
- [5] C. Breuil: *Cohomologie étale de p -torsion et cohomologie cristalline en réduction semi-stable*, Duke Math. J. **95** (1998), no. 3, 523–620.
- [6] C. Breuil: *Représentations semi-stables et modules fortement divisibles*, Invent. Math. **136** (1999), 89–122.
- [7] C. Breuil and W. Messing: *Torsion étale and crystalline cohomologies*, in *Cohomologies p -adiques et applications arithmétiques II*, Astérisque **279** (2002), 81–124.
- [8] X. Caruso: *Conjecture de l’inertie modérée de Serre*, Invent. Math. **171** (2008), 629–699.
- [9] X. Caruso and T. Liu: *Quasi-semi-stable representations*, to appear in Bull. Soc. Math. France.
- [10] J.-M. Fontaine: *Il n’y a pas de variété abélienne sur \mathbb{Z}* , Invent. Math. **81** (1985), 515–538.

- [11] J.-M. Fontaine: *Schémas propres et lisses sur \mathbb{Z}* , in *Proceedings of the Indo-French Conference on Geometry (Bombay, 1989)*, 43–56, 1993.
- [12] J.-M. Fontaine: *Le corps des périodes p -adiques*, *Astérisque* **223** (1994), 59–111.
- [13] J.-M. Fontaine and G. Laffaille: *Construction de représentations p -adiques*, *Ann. Sci. École Norm. Sup. (4)* **15** (1982) no. 4, 547–608.
- [14] J.-M. Fontaine and W. Messing: *p -adic periods and p -adic étale cohomology*, in *Current trends in arithmetical algebraic geometry (Arcata, Calif., 1985)*, *Contemp. Math.*, **67** (1987), 179–207.
- [15] M. Kisin: *Moduli of finite flat group schemes and modularity*, to appear in *Ann. of Math.*
- [16] M. Kisin: *Crystalline representations and F -crystals*, in *Algebraic Geometry and Number Theory*, *Progr. Math.* **253** (2006), 459–496.
- [17] T. Liu: *On lattices in semi-stable representations: a proof of a conjecture of Breuil*, *Compos. Math.* **144** (2008), 61–88.
- [18] J.-P. Serre: *Corps Locaux*, Hermann, Paris, 1968.
- [19] M. Yoshida: *Ramification of local fields and Fontaine’s property (P_m)* , preprint.
E-mail address: shin-h@math.kyushu-u.ac.jp

FACULTY OF MATHEMATICS, KYUSHU UNIVERSITY